

Myosin VI Polyclonal Antibody

Catalog # AP73255

Specification

Myosin VI Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9UM54
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Myosin VI Polyclonal Antibody - Additional Information

Gene ID 4646

Other Names

MYO6; KIAA0389; Unconventional myosin-VI; Unconventional myosin-6

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/5000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Myosin VI Polyclonal Antibody - Protein Information

Name MYO6 ([HGNC:7605](#))

Synonyms KIAA0389

Function

Myosins are actin-based motor molecules with ATPase activity (By similarity). Unconventional myosins serve in intracellular movements (By similarity). Myosin 6 is a reverse-direction motor protein that moves towards the minus-end of actin filaments (PubMed:10519557). Has slow rate of actin-activated ADP release due to weak ATP binding (By similarity). Functions in a variety of intracellular processes such as vesicular membrane trafficking and cell migration (By similarity). Required for the structural integrity of the Golgi apparatus via the p53-dependent pro-survival pathway (PubMed:16507995). Appears to be involved in a very early step of clathrin-mediated endocytosis in polarized epithelial cells (PubMed:11447109). Together with TOM1, mediates delivery of endocytic cargo to autophagosomes thereby promoting

autophagosome maturation and driving fusion with lysosomes (PubMed:23023224). Links TOM1 with autophagy receptors, such as TAX1BP1; CALCOCO2/NDP52 and OPTN (PubMed:31371777). May act as a regulator of F-actin dynamics (By similarity). As part of the DISP complex, may regulate the association of septins with actin and thereby regulate the actin cytoskeleton (PubMed:29467281). May play a role in transporting DAB2 from the plasma membrane to specific cellular targets (By similarity). May play a role in the extension and network organization of neurites (By similarity). Required for structural integrity of inner ear hair cells (By similarity). Required for the correct localization of CLIC5 and RDX at the stereocilium base (By similarity). Modulates RNA polymerase II- dependent transcription (PubMed:16949370).

Cellular Location

Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein. Golgi apparatus. Nucleus. Cytoplasm, perinuclear region. Membrane, clathrin-coated pit. Cytoplasmic vesicle, clathrin-coated vesicle. Cell projection, filopodium. Cell projection, ruffle membrane. Cell projection, microvillus. Cytoplasm, cytosol. Cytoplasmic vesicle, autophagosome. Endosome Note=Also present in endocytic vesicles (PubMed:16507995) Translocates from membrane ruffles, endocytic vesicles and cytoplasm to Golgi apparatus, perinuclear membrane and nucleus through induction by p53 and p53-induced DNA damage (PubMed:16507995). Recruited into membrane ruffles from cell surface by EGF-stimulation (PubMed:9852149) Colocalizes with DAB2 in clathrin-coated pits/vesicles (PubMed:11967127). Colocalizes with OPTN at the Golgi complex and in vesicular structures close to the plasma membrane (By similarity) Recruited to endosomes by TOM1 and TOM1L2 (PubMed:23023224) {ECO:0000250|UniProtKB:Q9I8D1, ECO:0000269|PubMed:11967127, ECO:0000269|PubMed:16507995, ECO:0000269|PubMed:23023224, ECO:0000269|PubMed:9852149} [Isoform 4]: Cytoplasmic vesicle, clathrin-coated vesicle membrane. Cell projection, ruffle membrane

Tissue Location




Expressed in most tissues examined including heart, brain, placenta, pancreas, spleen, thymus, prostate, testis, ovary, small intestine and colon. Highest levels in brain, pancreas, testis and small intestine. Also expressed in fetal brain and cochlea. Isoform 1 and isoform 2, containing the small insert, and isoform 4, containing neither insert, are expressed in unpolarized epithelial cells


Myosin VI Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Myosin VI Polyclonal Antibody - Images





Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100

Myosin VI Polyclonal Antibody - Background

Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Myosin 6 is a reverse-direction motor protein that moves towards the minus-end of actin filaments. Has slow rate of actin-activated ADP release due to weak ATP binding. Functions in a variety of intracellular processes such as vesicular membrane trafficking and cell migration. Required for the structural integrity of the Golgi apparatus via the p53-dependent pro-survival pathway. Appears to be involved in a very early step of clathrin-mediated endocytosis in polarized epithelial cells. May act as a regulator of F-actin dynamics. May play a role in transporting DAB2 from the plasma membrane to specific cellular targets. Required for structural integrity of inner ear hair cells (By similarity).