

TudorSN Polyclonal Antibody
Catalog # AP73267**Specification**

TudorSN Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q7KZF4
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

TudorSN Polyclonal Antibody - Additional Information**Gene ID** 27044**Other Names**

SND1; TDRD11; Staphylococcal nuclease domain-containing protein 1; 100 kDa coactivator; EBNA2 coactivator p100; Tudor domain-containing protein 11; p100 co-activator

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

TudorSN Polyclonal Antibody - Protein Information**Name** SND1**Synonyms** TDRD11**Function**

Endonuclease that mediates miRNA decay of both protein-free and AGO2-loaded miRNAs (PubMed:18453631, PubMed:28546213). As part of its function in miRNA decay, regulates mRNAs involved in G1-to-S phase transition (PubMed:28546213). Functions as a bridging factor between STAT6 and the basal transcription factor (PubMed:12234934). Plays a role in PIM1 regulation of MYB activity (PubMed:9809063). Functions as a transcriptional coactivator for STAT5 (By similarity).

Cellular Location

Cytoplasm. Nucleus. Melanosome Note=In IL-4 stimulated cells colocalizes with STAT6 in the nucleus (PubMed:12234934). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065)


Tissue Location

Ubiquitously expressed.


TudorSN Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TudorSN Polyclonal Antibody - Images


Western Blot analysis of extracts from Jurkat cells, using TudorSN Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100

TudorSN Polyclonal Antibody - Background

Endonuclease that mediates miRNA decay of both protein- free and AGO2-loaded miRNAs (PubMed:28546213, PubMed:18453631). As part of its function in miRNA decay, regulates mRNAs involved in G1-to-S phase transition (PubMed:28546213). Functions as a bridging factor between STAT6 and the basal transcription factor (PubMed:12234934). Plays a role in PIM1 regulation of MYB activity (PubMed:9809063). Functions as a transcriptional coactivator for STAT5 (By similarity).