

**SCML2 Polyclonal Antibody**  
**Catalog # AP73284****Specification**

---

**SCML2 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q9UQR0</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

**SCML2 Polyclonal Antibody - Additional Information****Gene ID** 10389**Other Names**

SCML2; Sex comb on midleg-like protein 2

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**SCML2 Polyclonal Antibody - Protein Information****Name** SCML2**Function**

Putative Polycomb group (PcG) protein. PcG proteins act by forming multiprotein complexes, which are required to maintain the transcriptionally repressive state of homeotic genes throughout development (By similarity).

**Cellular Location**

Nucleus.

**Tissue Location**



Highly expressed in placenta, thymus and testis. Detected at lower levels in brain, liver, skeletal muscle, pancreas and ovary.

**SCML2 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### SCML2 Polyclonal Antibody - Images



#### SCML2 Polyclonal Antibody - Background

Putative Polycomb group (PcG) protein. PcG proteins act by forming multiprotein complexes, which are required to maintain the transcriptionally repressive state of homeotic genes throughout development (By similarity).