

TRP14 Polyclonal Antibody
Catalog # AP73314**Specification**

TRP14 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q9BRA2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

TRP14 Polyclonal Antibody - Additional Information**Gene ID** 84817**Other Names**

TXNDC17; TXNL5; Thioredoxin domain-containing protein 17; 14 kDa thioredoxin-related protein; TRP14; Protein 42-9-9; Thioredoxin-like protein 5

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

TRP14 Polyclonal Antibody - Protein Information**Name** TXNDC17**Synonyms** TXNL5**Function**

Disulfide reductase. May participate in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions. Modulates TNF-alpha signaling and NF-kappa-B activation. Has peroxidase activity and may contribute to the elimination of cellular hydrogen peroxide.

Cellular Location

Cytoplasm.

Tissue Location



Ubiquitously expressed in cell lines.

TRP14 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRP14 Polyclonal Antibody - Images



TRP14 Polyclonal Antibody - Background

Disulfide reductase. May participate in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions. Modulates TNF-alpha signaling and NF-kappa-B activation. Has peroxidase activity and may contribute to the elimination of cellular hydrogen peroxide.