

CacyBP Polyclonal Antibody
Catalog # AP73572**Specification**

CacyBP Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9HB71
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

CacyBP Polyclonal Antibody - Additional Information**Gene ID** 27101**Other Names**

CACYPB; S100A6BP; SIP; PNAS-107; Calcyclin-binding protein; CacyBP; hCacyBP; S100A6-binding protein; Siah-interacting protein

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CacyBP Polyclonal Antibody - Protein Information**Name** CACYBP**Synonyms** S100A6BP, SIP**Function**

May be involved in calcium-dependent ubiquitination and subsequent proteasomal degradation of target proteins. Probably serves as a molecular bridge in ubiquitin E3 complexes. Participates in the ubiquitin-mediated degradation of beta-catenin (CTNNB1).

Cellular Location


Nucleus. Cytoplasm. Note=Cytoplasmic at low calcium concentrations. In neuroblastoma cells, after a retinoic acid (RA) induction and calcium increase, it localizes in both the nucleus and cytoplasm. The nuclear fraction may be phosphorylated

CacyBP Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CacyBP Polyclonal Antibody - Images




Western Blot analysis of HuvEc cells using CacyBP Polyclonal Antibody.. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-mammary-cancer, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-mammary-cancer, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100




Western Blot analysis of HuvEc cells using CacyBP Polyclonal Antibody.. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-mammary-cancer, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-mammary-cancer, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100

CacyBP Polyclonal Antibody - Background

May be involved in calcium-dependent ubiquitination and subsequent proteasomal degradation of target proteins. Probably serves as a molecular bridge in ubiquitin E3 complexes. Participates in the ubiquitin-mediated degradation of beta-catenin (CTNNB1).