

Eotaxin Polyclonal Antibody
Catalog # AP73627**Specification**

Eotaxin Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P51671
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Eotaxin Polyclonal Antibody - Additional Information**Gene ID** 6356**Other Names**

CCL11; SCYA11; Eotaxin; C-C motif chemokine 11; Eosinophil chemotactic protein; Small-inducible cytokine A11

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Eotaxin Polyclonal Antibody - Protein Information**Name** CCL11**Synonyms** SCYA11**Function**

In response to the presence of allergens, this protein directly promotes the accumulation of eosinophils, a prominent feature of allergic inflammatory reactions (PubMed:8597956). Binds to CCR3 (PubMed:8631813).

Cellular Location


Secreted.

Eotaxin Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Eotaxin Polyclonal Antibody - Images



Western Blot analysis of HepG2, mouse thymus, mouse skeletal muscle cells using Eotaxin Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of HepG2, mouse thymus, mouse skeletal muscle cells using Eotaxin Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Eotaxin Polyclonal Antibody - Background

In response to the presence of allergens, this protein directly promotes the accumulation of eosinophils, a prominent feature of allergic inflammatory reactions. Binds to CCR3.