

## FOXP3 Polyclonal Antibody

Catalog # AP73629

### Specification

#### FOXP3 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">Q9BZS1</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

#### FOXP3 Polyclonal Antibody - Additional Information

**Gene ID** 50943

#### Other Names

FOXP3; IPEX; JM2; Forkhead box protein P3; Scurfin

#### Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

#### Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

#### Storage Conditions

-20°C

#### FOXP3 Polyclonal Antibody - Protein Information

**Name** FOXP3

**Synonyms** IPEX

#### Function

Transcriptional regulator which is crucial for the development and inhibitory function of regulatory T-cells (Treg) (PubMed:<a href="http://www.uniprot.org/citations/17377532" target="\_blank">17377532</a>, PubMed:<a href="http://www.uniprot.org/citations/21458306" target="\_blank">21458306</a>, PubMed:<a href="http://www.uniprot.org/citations/23947341" target="\_blank">23947341</a>, PubMed:<a href="http://www.uniprot.org/citations/24354325" target="\_blank">24354325</a>, PubMed:<a href="http://www.uniprot.org/citations/24722479" target="\_blank">24722479</a>, PubMed:<a href="http://www.uniprot.org/citations/24835996" target="\_blank">24835996</a>, PubMed:<a href="http://www.uniprot.org/citations/30513302" target="\_blank">30513302</a>, PubMed:<a href="http://www.uniprot.org/citations/32644293" target="\_blank">32644293</a>). Plays an essential role in maintaining homeostasis of the immune system by allowing the acquisition of full suppressive function and stability of the Treg lineage, and by directly modulating the expansion and function of conventional T-cells (PubMed:<a href="http://www.uniprot.org/citations/32644293" target="\_blank">32644293</a>).

href="http://www.uniprot.org/citations/23169781" target="\_blank">23169781). Can act either as a transcriptional repressor or a transcriptional activator depending on its interactions with other transcription factors, histone acetylases and deacetylases (PubMed:<a href="http://www.uniprot.org/citations/17377532" target="\_blank">173775322145830623947341243543252472247917377532214583062394734124354325247224791579068117360565173775321836804918354202

### Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00089, ECO:0000269|PubMed:17360565,

ECO:0000269|PubMed:18354202, ECO:0000269|PubMed:22678915,

ECO:0000269|PubMed:23396208, ECO:0000269|PubMed:23973222,

ECO:0000269|PubMed:23973223, ECO:0000269|PubMed:32644293}. Cytoplasm


Note=Predominantly expressed in the cytoplasm in activated conventional T-cells whereas predominantly expressed in the nucleus in regulatory T- cells (Treg). The 41 kDa form derived by proteolytic processing is found exclusively in the chromatin fraction of activated Treg cells (By similarity). {ECO:0000250|UniProtKB:Q99JB6, ECO:0000269|PubMed:22678915}

### FOXP3 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### FOXP3 Polyclonal Antibody - Images




Western Blot analysis of PC12 cells using FOXP3 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-tonsilla, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-tonsilla, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100

#### **FOXP3 Polyclonal Antibody - Background**

Transcriptional regulator which is crucial for the development and inhibitory function of regulatory T-cells (Treg). Plays an essential role in maintaining homeostasis of the immune system by allowing the acquisition of full suppressive function and stability of the Treg lineage, and by directly modulating the expansion and function of conventional T-cells. Can act either as a transcriptional repressor or a transcriptional activator depending on its interactions with other transcription factors, histone acetylases and deacetylases. The suppressive activity of Treg involves the coordinate activation of many genes, including CTLA4 and TNFRSF18 by FOXP3 along with repression of genes encoding cytokines such as interleukin-2 (IL2) and interferon- gamma (IFNG). Inhibits cytokine production and T-cell effector function by repressing the activity of two key transcription factors, RELA and NFATC2 (PubMed:15790681). Mediates transcriptional repression of IL2 via its association with histone acetylase KAT5 and histone deacetylase HDAC7

(PubMed:17360565). Can activate the expression of TNFRSF18, IL2RA and CTLA4 and repress the expression of IL2 and IFNG via its association with transcription factor RUNX1 (PubMed:17377532). Inhibits the differentiation of IL17 producing helper T-cells (Th17) by antagonizing RORC function, leading to down-regulation of IL17 expression, favoring Treg development (PubMed:18368049). Inhibits the transcriptional activator activity of RORA (PubMed:18354202). Can repress the expression of IL2 and IFNG via its association with transcription factor IKZF4 (By similarity).