

**LIR-7 Polyclonal Antibody**  
**Catalog # AP73649****Specification**

---

**LIR-7 Polyclonal Antibody - Product Information**

Application	<b>WB, IHC-P</b>
Primary Accession	<a href="#">Q8N149</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>

**LIR-7 Polyclonal Antibody - Additional Information****Gene ID** 11027**Other Names**

LILRA2; ILT1; LIR7; Leukocyte immunoglobulin-like receptor subfamily A member 2; CD85 antigen-like family member H; Immunoglobulin-like transcript 1; ILT-1; Leukocyte immunoglobulin-like receptor 7; LIR-7; CD85h

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.  
IHC-P~~N/A

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**LIR-7 Polyclonal Antibody - Protein Information****Name** LILRA2**Synonyms** ILT1, LIR7**Function**

Part of the innate immune responses against microbial infection (PubMed:<a href="http://www.uniprot.org/citations/12529506" target="\_blank">12529506</a>, PubMed:<a href="http://www.uniprot.org/citations/27572839" target="\_blank">27572839</a>). Specifically recognizes a set of N-terminally truncated immunoglobulins that are produced via cleavage by proteases from a range of pathogenic bacteria and fungi, including L.pneumophila, M.hyorhinis, S.pneumoniae, S.aureus and C.albicans (PubMed:<a href="http://www.uniprot.org/citations/27572839" target="\_blank">27572839</a>). Recognizes epitopes that are in part in the variable region of the immunoglobulin light chains, but requires also the constant region for signaling (PubMed:<a href="http://www.uniprot.org/citations/27572839" target="\_blank">27572839</a>). Binds to a

subset of cleaved IgM, IgG3 and IgG4 molecules, but does not bind cleaved IgA1 (PubMed:<a href="http://www.uniprot.org/citations/27572839" target="\_blank">27572839</a>). Binding of N-terminally truncated immunoglobulins mediates activation of neutrophils (PubMed:<a href="http://www.uniprot.org/citations/27572839" target="\_blank">27572839</a>). In monocytes, activation leads to the release of CSF2, CF3, IL6, CXCL8 and CCL3 and down-regulates responses to bacterial lipopolysaccharide (LPS), possibly via down-regulation of TLR4 expression and reduced signaling via TLR4 (PubMed:<a href="http://www.uniprot.org/citations/22479404" target="\_blank">22479404</a>). In eosinophils, activation by ligand binding leads to the release of RNASE2, IL4 and leukotriene C4 (PubMed:<a href="http://www.uniprot.org/citations/12529506" target="\_blank">12529506</a>). Does not bind class I MHC antigens (PubMed:<a href="http://www.uniprot.org/citations/19230061" target="\_blank">19230061</a>).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein

#### Tissue Location


Detected on the surface of all peripheral blood monocytes, neutrophils, basophils and eosinophils (at protein level) (PubMed:12529506, PubMed:22479404). Expression levels are very low or not detectable on monocytes, T-cells, B-cells, dendritic cells and natural killer (NK) cells (PubMed:9548455)





#### LIR-7 Polyclonal Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.



- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### LIR-7 Polyclonal Antibody - Images









### **LIR-7 Polyclonal Antibody - Background**

Part of the innate immune responses against microbial infection (PubMed:12529506, PubMed:27572839). Specifically recognizes a set of N-terminally truncated immunoglobulins that are produced via cleavage by proteases from a range of pathogenic bacteria and fungi, including *L.pneumophila*, *M.hyorhinis*, *S.pneumoniae*, *S.aureus* and *C.albicans* (PubMed:27572839). Recognizes epitopes that are in part in the variable region of the immunoglobulin light chains, but requires also the constant region for signaling (PubMed:27572839). Binds to a subset of cleaved IgM, IgG3 and IgG4 molecules, but does not bind cleaved IgA1 (PubMed:27572839). Binding of N-terminally truncated immunoglobulins mediates activation of neutrophils (PubMed:27572839). In monocytes, activation leads to the release of CSF2, CF3, IL6, CXCL8 and CCL3 and down-regulates responses to bacterial lipopolysaccharide (LPS), possibly via down-regulation of TLR4 expression and reduced signaling via TLR4 (PubMed:22479404). In eosinophils, activation by ligand binding leads to the release of RNASE2, IL4 and leukotriene C4 (PubMed:12529506). Does not bind class I MHC antigens (PubMed:19230061).