

CD200 Polyclonal Antibody
Catalog # AP73657**Specification**

CD200 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P41217
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

CD200 Polyclonal Antibody - Additional Information**Gene ID** 4345**Other Names**

CD200; MOX1; MOX2; My033; OX-2 membrane glycoprotein; CD200

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CD200 Polyclonal Antibody - Protein Information**Name** CD200**Synonyms** MOX1, MOX2**Function**

Costimulates T-cell proliferation. May regulate myeloid cell activity in a variety of tissues.

Cellular Location

Cell membrane; Single-pass type I membrane protein


CD200 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CD200 Polyclonal Antibody - Images




Western Blot analysis of PC12 cells using CD200 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100

CD200 Polyclonal Antibody - Background

Costimulates T-cell proliferation. May regulate myeloid cell activity in a variety of tissues.