

BST-1 Polyclonal Antibody
Catalog # AP73659**Specification**

BST-1 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q10588
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

BST-1 Polyclonal Antibody - Additional Information**Gene ID** 683**Other Names**

BST1; ADP-ribosyl cyclase 2; Bone marrow stromal antigen 1; BST-1; Cyclic ADP-ribose hydrolase 2; cADPr hydrolase 2; CD157

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

BST-1 Polyclonal Antibody - Protein Information**Name** BST1**Function**

Catalyzes both the synthesis of cyclic ADP-beta-D-ribose (cADPR) from NAD(+), and its hydrolysis to ADP-D-ribose (ADPR) (PubMed:7805847). Cyclic ADPR is known to serve as an endogenous second messenger that elicits calcium release from intracellular stores, and thus regulates the mobilization of intracellular calcium (Probable). May be involved in pre-B-cell growth (Probable).

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor

Tissue Location



Expressed in various tissues including placenta, lung, liver and kidney.

BST-1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BST-1 Polyclonal Antibody - Images



BST-1 Polyclonal Antibody - Background

Synthesizes the second messengers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger that elicits calcium release from intracellular stores. May be involved in pre-B-cell growth.