

NKp44 Polyclonal Antibody
Catalog # AP73664**Specification**

NKp44 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	O95944
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

NKp44 Polyclonal Antibody - Additional Information**Gene ID** 9436**Other Names**

NCR2; LY95; Natural cytotoxicity triggering receptor 2; Lymphocyte antigen 95 homolog; NK cell-activating receptor; Natural killer cell p44-related protein; NK-p44; NKp44; CD336

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

NKp44 Polyclonal Antibody - Protein Information**Name** NCR2**Synonyms** LY95**Function**

Cytotoxicity-activating receptor that may contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location



Selectively expressed by activated NK cells and by in vitro cultured (i.e. activated) TCRg/d lymphoid cells




NKp44 Polyclonal Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NKp44 Polyclonal Antibody - Images







NKp44 Polyclonal Antibody - Background

Cytotoxicity-activating receptor that may contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis.