

ICAM-2 Polyclonal Antibody

Catalog # AP73784

Specification

ICAM-2 Polyclonal Antibody - Product Information

Application WB
Primary Accession P13598
Reactivity Human
Host Rabbit
Clonality Polyclonal

ICAM-2 Polyclonal Antibody - Additional Information

Gene ID 3384

Other Names

ICAM2; Intercellular adhesion molecule 2; ICAM-2; CD102

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

ICAM-2 Polyclonal Antibody - Protein Information

Name ICAM2

Function

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance.

Cellular Location

Membrane; Single-pass type I membrane protein. Cell projection, microvillus {ECO:0000250|UniProtKB:P35330}. Note=Co-localizes with RDX, EZR and MSN in microvilli. {ECO:0000250|UniProtKB:P35330}

ICAM-2 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.





- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

ICAM-2 Polyclonal Antibody - Images



ICAM-2 Polyclonal Antibody - Background

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen- specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance.