

GDF-7 Polyclonal Antibody
Catalog # AP73837**Specification**

GDF-7 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q7Z4P5
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

GDF-7 Polyclonal Antibody - Additional Information**Gene ID** 151449**Other Names**

GDF7; Growth/differentiation factor 7; GDF-7

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

GDF-7 Polyclonal Antibody - Protein Information**Name** GDF7**Function**

May play an active role in the motor area of the primate neocortex.

Cellular Location

Secreted.


GDF-7 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


GDF-7 Polyclonal Antibody - Images




Western Blot analysis of Hela cells using GDF-7 Polyclonal Antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Western Blot analysis of Hela cells using GDF-7 Polyclonal Antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

GDF-7 Polyclonal Antibody - Background

May play an active role in the motor area of the primate neocortex.