

Factor XIII B Polyclonal Antibody
Catalog # AP73884**Specification**

Factor XIII B Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P05160
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Factor XIII B Polyclonal Antibody - Additional Information**Gene ID** 2165**Other Names**

F13B; Coagulation factor XIII B chain; Fibrin-stabilizing factor B subunit; Protein-glutamine gamma-glutamyltransferase B chain; Transglutaminase B chain

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Factor XIII B Polyclonal Antibody - Protein Information**Name** F13B**Function**

The B chain of factor XIII is not catalytically active, but is thought to stabilize the A subunits and regulate the rate of transglutaminase formation by thrombin.

Cellular Location

Secreted


Factor XIII B Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Factor XIII B Polyclonal Antibody - Images



Western blot analysis of various lysates using F13B antibody. Antibody was diluted at 1:2000. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human liver, antibody was diluted at 1:200

Factor XIII B Polyclonal Antibody - Background

The B chain of factor XIII is not catalytically active, but is thought to stabilize the A subunits and regulate the rate of transglutaminase formation by thrombin.