

**IMPDH2 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7390c****Specification**

---

**IMPDH2 Antibody (Center) - Product Information**

Application	FC, IHC-P, WB,E
Primary Accession	<a href="#">P12268</a>
Other Accession	<a href="#">E9PU28</a> , <a href="#">P24547</a> , <a href="#">B0UXP9</a> , <a href="#">P12269</a> , <a href="#">Q3SWY3</a> , <a href="#">D3ZLZ7</a> , <a href="#">P50096</a> , <a href="#">P20839</a> , <a href="#">A0JNA3</a>
Reactivity	Human
Predicted	Bovine, Mouse, Rat, Hamster, Zebrafish
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	55805
Antigen Region	180-210

**IMPDH2 Antibody (Center) - Additional Information****Gene ID** 3615**Other Names**

Inosine-5'-monophosphate dehydrogenase 2 {ECO:0000255|HAMAP-Rule:MF\_03156}, IMP dehydrogenase 2 {ECO:0000255|HAMAP-Rule:MF\_03156}, IMPD 2 {ECO:0000255|HAMAP-Rule:MF\_03156}, IMPDH 2 {ECO:0000255|HAMAP-Rule:MF\_03156}, 111205 {ECO:0000255|HAMAP-Rule:MF\_03156}, IMPDH-II, IMPDH2 {ECO:0000255|HAMAP-Rule:MF\_03156}, IMPD2

**Target/Specificity**

This IMPDH2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 180-210 amino acids from the Central region of human IMPDH2.

**Dilution**

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

IMPDH2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## IMPDH2 Antibody (Center) - Protein Information

**Name** IMPDH2 ([HGNC:6053](#))

**Synonyms** IMPD2

**Function** Catalyzes the conversion of inosine 5'-phosphate (IMP) to xanthosine 5'-phosphate (XMP), the first committed and rate-limiting step in the de novo synthesis of guanine nucleotides, and therefore plays an important role in the regulation of cell growth (PubMed:[7763314](#), PubMed:[7903306](#)). Could also have a single-stranded nucleic acid-binding activity and could play a role in RNA and/or DNA metabolism (PubMed:[14766016](#)). It may also have a role in the development of malignancy and the growth progression of some tumors.

### Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytosol. Note=Can form fiber-like subcellular structures termed 'cytoophidia' in response to intracellular guanine- nucleotide depletion.

### Tissue Location


IMPDH1 is the main species in normal leukocytes and IMPDH2 predominates over IMPDH1 in the tumor

## IMPDH2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## IMPDH2 Antibody (Center) - Images




Western blot analysis of IMPDH2 Antibody (Center) (Cat. #AP7390c) in NCI-H460, HeLa, MCF7, A2058 cell line lysates (35ug/lane). IMPDH2 (arrow) was detected using the

purified Pab.



IMPDH2 Antibody (Center) (Cat. #AP7390c) IHC analysis in formalin fixed and paraffin embedded human Lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the IMPDH2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



IMPDH2 Antibody (Center) (Cat. #AP7390c) flow cytometric analysis of MCF-7 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### IMPDH2 Antibody (Center) - Background

IMPDH2 is the rate-limiting enzyme in the de novo guanine nucleotide biosynthesis. It is thus involved in maintaining cellular guanine deoxy- and ribonucleotide pools needed for DNA and RNA synthesis. The protein catalyzes the NAD-dependent oxidation of inosine-5'-monophosphate into xanthine-5'-monophosphate, which is then converted into guanosine-5'-monophosphate. Its gene is up-regulated in some neoplasms, suggesting it may play a role in malignant transformation.

### IMPDH2 Antibody (Center) - References

Sombogaard, F., Pharmacogenet. Genomics 19 (8), 626-634 (2009)  
 Mohamed, M.F., Genet. Test. 12 (4), 513-516 (2008)