

C32 Polyclonal Antibody
Catalog # AP73913**Specification**

C32 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q9UBR5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

C32 Polyclonal Antibody - Additional Information**Gene ID** 51192**Other Names**

CKLF; CKLF1; HSPC224; Chemokine-like factor; C32

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

C32 Polyclonal Antibody - Protein Information**Name** CKLF**Synonyms** CKLF1**Function**

May play an important role in inflammation and regeneration of skeletal muscle (PubMed:11415443). Essential for embryonic development (By similarity).

Cellular Location

[Isoform 1]: Secreted [Isoform 4]: Membrane; Multi-pass membrane protein

Tissue Location



Isoform 1, isoform 2, isoform 3 and isoform 4 have highest expression levels in adult spleen, lung, testis, ovary, peripheral blood leukocyte, placenta, pancreas, and in fetal brain, skeletal muscle, thymus and heart. Lower expression levels in adult skeletal muscle, liver, thymus colon, prostate and fetal spleen and liver.

C32 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C32 Polyclonal Antibody - Images



C32 Polyclonal Antibody - Background

May play an important role in inflammation and regeneration of skeletal muscle. Partly inhibited

by interleukin 10.