

TGFβ RIII Polyclonal Antibody
Catalog # AP73938**Specification**

TGFβ RIII Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q03167
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

TGFβ RIII Polyclonal Antibody - Additional Information**Gene ID** 7049**Other Names**

transforming growth factor, beta receptor III

Dilution

WB~~WB 1:500-2000, ELISA 1:10000-20000

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

TGFβ RIII Polyclonal Antibody - Protein Information**Name** TGFBR3 ([HGNC:11774](#))**Function**

Cell surface receptor that regulates diverse cellular processes including cell proliferation, differentiation, migration, and apoptosis (PubMed:12958365, PubMed:19416857). Initiates BMP, inhibin, and TGF-beta signaling pathways by interacting with different ligands including TGFβ1, BMP2, BMP5, BMP7 or GDF5 (PubMed:18184661). Alternatively, acts as a cell surface coreceptor for BMP ligands, serving to enhance ligand binding by differentially regulating BMPRI/ALK3 and BMPRII/ALK6 receptor trafficking (PubMed:19726563). Promotes epithelial cell adhesion, focal adhesion formation and integrin signaling during epithelial cell spreading on fibronectin (PubMed:22562249). By interacting with the scaffolding protein beta-arrestin2/ARRB2, regulates migration or actin cytoskeleton and promotes the activation of CDC42 as well as the inhibition of NF-kappa-B (PubMed:19416857, PubMed:19416857).

href="http://www.uniprot.org/citations/19325136" target="_blank">19325136). In gonadotrope cells, acts as an inhibin A coreceptor and regulates follicle-stimulating hormone (FSH) levels and female fertility (By similarity). Plays a role in the inhibition of directed and random cell migration in epithelial cells by altering the actin cytoskeletal organization (PubMed:19416857). Participates in epithelial-mesenchymal transformation (EMT) upon binding to BMP2 or TGFB2, by activating the PAR6/SMURF1/RHOA pathway (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Secreted {ECO:0000250|UniProtKB:P26342}. Secreted, extracellular space, extracellular matrix {ECO:0000250|UniProtKB:P26342}. Note=Exists both as a membrane-bound form and as soluble form in serum and in the extracellular matrix. {ECO:0000250|UniProtKB:P26342}

TGFβ RIII Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TGFβ RIII Polyclonal Antibody - Images





Image not found : 202004/c30783wb42457.jpg



TGFβ RIII Polyclonal Antibody - Background

Binds to TGF-beta. Could be involved in capturing and retaining TGF-beta for presentation to the signaling receptors.