

CD204 Polyclonal Antibody

Catalog # AP74130

Specification

CD204 Polyclonal Antibody - Product Information

Application IHC-P Primary Accession P21757

Reactivity Human, Mouse, Rat

Host Rabbit Clonality Polyclonal

CD204 Polyclonal Antibody - Additional Information

Gene ID 4481

Other Names

Macrophage scavenger receptor types I and II (Macrophage acetylated LDL receptor I and II) (Scavenger receptor class A member 1) (CD antigen CD204)

Dilution

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CD204 Polyclonal Antibody - Protein Information

Name MSR1

Synonyms SCARA1

Function

Membrane glycoproteins implicated in the pathologic deposition of cholesterol in arterial walls during atherogenesis. Two types of receptor subunits exist. These receptors mediate the endocytosis of a diverse group of macromolecules, including modified low density lipoproteins (LDL) (PubMed:<a href="http://www.uniprot.org/citations/2251254"

target="_blank">2251254). Isoform III does not internalize acetylated LDL (PubMed:9548586).

Cellular Location

Membrane; Single-pass type II membrane protein.

Tissue Location

Isoform I, isoform II and isoform III are expressed in monocyte-derived macrophages. Isoform I and isoform II are expressed in the liver, placenta and brain.




CD204 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CD204 Polyclonal Antibody - Images




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200





Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200

CD204 Polyclonal Antibody - Background

Membrane glycoproteins implicated in the pathologic deposition of cholesterol in arterial walls during atherogenesis. Two types of receptor subunits exist. These receptors mediate the endocytosis of a diverse group of macromolecules, including modified low density lipoproteins (LDL) (PubMed:2251254). Isoform III does not internalize acetylated LDL (PubMed:9548586).