

NNT-1 Polyclonal Antibody
Catalog # AP74140**Specification****NNT-1 Polyclonal Antibody - Product Information**

| | |
|-------------------|------------------------|
| Application | IHC-P |
| Primary Accession | Q9UBD9 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |

NNT-1 Polyclonal Antibody - Additional Information**Gene ID** 23529**Other Names**

Cardiotrophin-like cytokine factor 1 (B-cell-stimulating factor 3) (BSF-3) (Novel neurotrophin-1) (NNT-1)

Dilution

IHC-P~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

NNT-1 Polyclonal Antibody - Protein Information**Name** CLCF1**Synonyms** BSF3, CLC, NNT1**Function**

In complex with CRLF1, forms a heterodimeric neurotropic cytokine that plays a crucial role during neuronal development (Probable). Also stimulates B-cells. Binds to and activates the ILST/gp130 receptor.

Cellular Location

Secreted.

Tissue Location


Expressed predominantly in lymph nodes, spleen, peripheral blood lymphocytes, bone marrow, and fetal liver

NNT-1 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


NNT-1 Polyclonal Antibody - Images




Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

NNT-1 Polyclonal Antibody - Background

Cytokine with B-cell stimulating capability. Binds to and activates the ILST/gp130 receptor.