

Eotaxin-2 Polyclonal Antibody
Catalog # AP74142**Specification**

Eotaxin-2 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O00175
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Eotaxin-2 Polyclonal Antibody - Additional Information**Gene ID** 6369**Other Names**

C-C motif chemokine 24 (CK-beta-6) (Eosinophil chemotactic protein 2) (Eotaxin-2) (Myeloid progenitor inhibitory factor 2) (MPIF-2) (Small-inducible cytokine A24)

Dilution

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Eotaxin-2 Polyclonal Antibody - Protein Information**Name** CCL24 ([HGNC:10623](#))**Function**

Chemotactic for resting T-lymphocytes, and eosinophils (PubMed:9104803, PubMed:9365122). Has lower chemotactic activity for neutrophils but none for monocytes and activated lymphocytes (PubMed:9104803, PubMed:9365122). Is a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line (PubMed:9104803, PubMed:9365122). Binds to CCR3 (PubMed:9104803, PubMed:9365122).

Cellular Location

Secreted


Tissue Location

Activated monocytes and activated T lymphocytes.


Eotaxin-2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Eotaxin-2 Polyclonal Antibody - Images


Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:200

Eotaxin-2 Polyclonal Antibody - Background

Chemotactic for resting T-lymphocytes, and eosinophils. Has lower chemotactic activity for neutrophils but none for monocytes and activated lymphocytes. Is a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line. Binds to CCR3.