

GRO γ Polyclonal Antibody
Catalog # AP74153**Specification**

GRO γ Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P19876
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

GRO γ Polyclonal Antibody - Additional Information**Gene ID** 2921**Other Names**

C-X-C motif chemokine 3 (GRO-gamma(1-73)) (Growth-regulated protein gamma) (GRO-gamma) (Macrophage inflammatory protein 2-beta) (MIP2-beta) [Cleaved into: GRO-gamma(5-73)]

Dilution

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

GRO γ Polyclonal Antibody - Protein Information**Name** CXCL3**Synonyms** GRO3, GROG, SCYB3**Function**

Ligand for CXCR2 (By similarity). Has chemotactic activity for neutrophils. May play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion. In vitro, the processed form GRO-gamma(5-73) shows a fivefold higher chemotactic activity for neutrophilic granulocytes.

Cellular Location



Secreted.

GRO γ Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GROγ Polyclonal Antibody - Images



GROγ Polyclonal Antibody - Background

Ligand for CXCR2 (By similarity). Has chemotactic activity for neutrophils. May play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion. In vitro, the processed form GRO-gamma(5-73) shows a fivefold higher chemotactic activity for neutrophilic granulocytes.