

GSTA1 Polyclonal Antibody
Catalog # AP74164**Specification****GSTA1 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	P08263
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

GSTA1 Polyclonal Antibody - Additional Information**Gene ID** 2938**Other Names**

Glutathione S-transferase A1 (EC 2.5.1.18) (GST HA subunit 1) (GST class-alpha member 1) (GST-epsilon) (GSTA1-1) (GTH1)

Dilution

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

GSTA1 Polyclonal Antibody - Protein Information**Name** GSTA1**Function**

Glutathione S-transferase that catalyzes the nucleophilic attack of the sulfur atom of glutathione on the electrophilic groups of a wide range of exogenous and endogenous compounds (Probable). Involved in the formation of glutathione conjugates of both prostaglandin A2 (PGA2) and prostaglandin J2 (PGJ2) (PubMed: [9084911](http://www.uniprot.org/citations/9084911) target="_blank">9084911). It also catalyzes the isomerization of D5-androstene-3,17-dione (AD) into D4-androstene- 3,17-dione and may therefore play an important role in hormone biosynthesis (PubMed: [11152686](http://www.uniprot.org/citations/11152686) target="_blank">11152686). Through its glutathione-dependent peroxidase activity toward the fatty acid hydroperoxide (13S)- hydroperoxy-(9Z,11E)-octadecadienoate/13-HPODE it is also involved in the metabolism of oxidized linoleic acid (PubMed: [16624487](http://www.uniprot.org/citations/16624487) target="_blank">16624487).

Cellular Location

Cytoplasm.

Tissue Location


Liver.

GSTA1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GSTA1 Polyclonal Antibody - Images



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200

GSTA1 Polyclonal Antibody - Background

Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles.