

MCP-3 Polyclonal Antibody
Catalog # AP74229**Specification****MCP-3 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	P80098
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

MCP-3 Polyclonal Antibody - Additional Information**Gene ID** 6354**Other Names**

C-C motif chemokine 7 (Monocyte chemoattractant protein 3) (Monocyte chemotactic protein 3) (MCP-3) (NC28) (Small-inducible cytokine A7)

Dilution

IHC-P~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

MCP-3 Polyclonal Antibody - Protein Information**Name** CCL7**Synonyms** MCP3, SCYA6, SCYA7**Function**

Chemotactic factor that attracts monocytes and eosinophils, but not neutrophils. Augments monocyte anti-tumor activity. Also induces the release of gelatinase B. This protein can bind heparin. Binds to CCR1, CCR2 and CCR3.


Cellular Location

Secreted.


MCP-3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MCP-3 Polyclonal Antibody - Images


Immunohistochemical analysis of paraffin-embedded Human-skin, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded Human-colon, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded Human-colon, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-skin, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-colon, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-colon, antibody was diluted at 1:100
MCP-3 Polyclonal Antibody - Background

Chemotactic factor that attracts monocytes and eosinophils, but not neutrophils. Augments monocyte anti-tumor activity. Also induces the release of gelatinase B. This protein can bind heparin. Binds to CCR1, CCR2 and CCR3.