

**FA2H Polyclonal Antibody**  
**Catalog # AP74261****Specification**

---

**FA2H Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q7L5A8</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**FA2H Polyclonal Antibody - Additional Information****Gene ID** 79152**Other Names**

Fatty acid 2-hydroxylase (EC 1.-.-.) (Fatty acid alpha-hydroxylase)

**Dilution**

WB~~WB 1:500-2000, ELISA 1:10000-20000

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**FA2H Polyclonal Antibody - Protein Information****Name** FA2H**Function**

Catalyzes the hydroxylation of free fatty acids at the C-2 position to produce 2-hydroxy fatty acids, which are building blocks of sphingolipids and glycosphingolipids common in neural tissue and epidermis (PubMed: [15337768](http://www.uniprot.org/citations/15337768), PubMed: [15863841](http://www.uniprot.org/citations/15863841), PubMed: [17355976](http://www.uniprot.org/citations/17355976), PubMed: [22517924](http://www.uniprot.org/citations/22517924)). FA2H is stereospecific for the production of (R)-2- hydroxy fatty acids (PubMed: [22517924](http://www.uniprot.org/citations/22517924)). Plays an essential role in the synthesis of galactosphingolipids of the myelin sheath (By similarity). Responsible for the synthesis of sphingolipids and glycosphingolipids involved in the formation of epidermal lamellar bodies critical for skin permeability barrier (PubMed: [17355976](http://www.uniprot.org/citations/17355976)). Participates in the synthesis of glycosphingolipids and a fraction of type II wax diesters in sebaceous gland, specifically regulating hair follicle homeostasis (By similarity). Involved in the synthesis of sphingolipids of plasma membrane rafts, controlling lipid raft mobility and trafficking of raft-associated proteins (By similarity).

**Cellular Location**

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q5MPP0}; Multi-pass membrane protein. Microsome membrane; Multi-pass membrane protein


**Tissue Location**


Detected in differentiating cultured keratinocytes (at protein level). Detected in epidermis and cultured keratinocytes (PubMed:17355976). Highly expressed in brain and colon. Detected at lower levels in testis, prostate, pancreas and kidney (PubMed:15337768).

**FA2H Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**FA2H Polyclonal Antibody - Images**



#### **FA2H Polyclonal Antibody - Background**

Catalyzes stereospecific hydroxylation of free fatty acids at the C-2 position to produce (R)-2-hydroxy fatty acids, which are building blocks of sphingolipids and glycosphingolipids common in neural tissue and epidermis (PubMed:17355976, PubMed:22517924, PubMed:15863841, PubMed:15337768). Plays an essential role in the synthesis of galactosphingolipids of the myelin sheath (By similarity). Responsible for the synthesis of sphingolipids and glycosphingolipids involved in the formation of epidermal lamellar bodies critical for skin permeability barrier (PubMed:17355976). Participates in the synthesis of glycosphingolipids and a fraction of type II wax diesters in sebaceous gland, specifically regulating hair follicle homeostasis (By similarity). Involved in the synthesis of sphingolipids of plasma membrane rafts, controlling lipid raft mobility and trafficking of raft-associated proteins (By similarity).