

FASN Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7449C**Specification**

FASN Antibody (Center) - Product Information

Application	FC, IHC-P, IF, WB,E
Primary Accession	P49327
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	273427
Antigen Region	942-973

FASN Antibody (Center) - Additional Information**Gene ID** 2194**Other Names**

Fatty acid synthase, [Acyl-carrier-protein] S-acetyltransferase, [Acyl-carrier-protein] S-malonyltransferase, 3-oxoacyl-[acyl-carrier-protein] synthase, 3-oxoacyl-[acyl-carrier-protein] reductase, 3-hydroxyacyl-[acyl-carrier-protein] dehydratase, Enoyl-[acyl-carrier-protein] reductase, Oleoyl-[acyl-carrier-protein] hydrolase, FASN, FAS

Target/Specificity

This FASN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 942-973 amino acids from the Central region of human FASN.

Dilution

FC~~1:10~50
IHC-P~~1:10~50
IF~~1:10~50
WB~~1:16000
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FASN Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

FASN Antibody (Center) - Protein Information

Name FASN

Synonyms FAS

Function Fatty acid synthetase is a multifunctional enzyme that catalyzes the de novo biosynthesis of long-chain saturated fatty acids starting from acetyl-CoA and malonyl-CoA in the presence of NADPH. This multifunctional protein contains 7 catalytic activities and a site for the binding of the prosthetic group 4'-phosphopantetheine of the acyl carrier protein ([ACP]) domain.

Cellular Location

Cytoplasm. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

Tissue Location


Ubiquitous. Prominent expression in brain, lung, liver and mammary gland.

FASN Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FASN Antibody (Center) - Images




Fluorescent confocal image of HepG2 cells stained with FASN (Center) antibody. HepG2 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AP7449c FASN (Center) primary antibody (1:200, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 µg/ml, 5 min). Note the


highly specific localization of the FASN mainly to the cytoplasm, supported by Human Protein Atlas Data (<http://www.proteinatlas.org/ENSG00000169710>).



Confocal immunofluorescent analysis of FASN Antibody (Center)(Cat#AP7449c) with HeLa cells followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclei (blue).




Western blot analysis of FASN Antibody (Center) (Cat. #AP7449c) in Ramos cell line lysates (35ug/lane). FASN (arrow) was detected using the purified Pab.




Western blot analysis of FASN (arrow) using rabbit polyclonal FASN Antibody


(Center)(Cat. #AP7449c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the FASN gene (Lane 2) (Origene Technologies).



All lanes : Anti-FASN Antibody (Center) at 1:16000 dilution Lane 1: A549 whole cell lysate Lane 2: Hela whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: SH-SY5Y whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 273 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



FASN Antibody (Center) (Cat. #AP7449c) immunohistochemistry analysis in formalin fixed and paraffin embedded human placenta tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of FASN Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



FASN Antibody (Center) (Cat. #AP7449c) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FASN Antibody (Center) - Background

FASN is a multifunctional protein. Its main function is to catalyze the synthesis of palmitate from acetyl-CoA and malonyl-CoA, in the presence of NADPH, into long-chain saturated fatty acids. In some cancer cell lines, this protein has been found to be fused with estrogen receptor-alpha (ER-alpha), in which the N-terminus of FAS is fused in-frame with the C-terminus of ER-alpha.

FASN Antibody (Center) - References

References for protein:

1. Jayakumar A., Tai M.-H. Proc. Natl. Acad. Sci. U.S.A. 92:8695-8699(1995)
2. Kuhajda F.P., Jenner K. Proc. Natl. Acad. Sci. U.S.A. 91:6379-6383(1994)
3. Semenkovich C.F., Coleman T.J. Lipid Res. 36:1507-1521(1995)

References for HepG2 cell line:

1. Knowles BB, et al. (1980). Human hepatocellular carcinoma cell lines secrete the major plasma proteins and hepatitis B surface antigen. Science 209: 497-499. [PubMed: 6248960].
2. Darlington GJ, et al. (1987). Growth and hepatospecific gene expression of human hepatoma cells in a defined medium. In Vitro Cell. Dev. Biol. 23: 349-354. [PubMed: 3034851].
3. Ihrke, G; Neufeld, EB; Meads, T; Shanks, MR; Cassio, D; Laurent, M; Schroer, TA; Pagano, RE et al. (1993). "WIF-B cells: an in vitro model for studies of hepatocyte polarity". Journal of Cell Biology 123 (6): 1761-1775. [PubMed:7506266].
4. Mersch-Sundermann, V.; Knasmüller, S.; Wu, X. J.; Darroudi, F.; Kassie, F. (2004). "Use of a human-derived liver cell line for the detection of cytoprotective, antigenotoxic and cogenotoxic agents". Toxicology 198 (1-3): 329-340. [PubMed:15138059].