

DDX39 Antibody (Center Y265)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7451d

Specification

DDX39 Antibody (Center Y265) - Product Information

Application WB,E
Primary Accession 000148

Other Accession Q5U216, Q8VDW0

Reactivity
Predicted
Mouse, Rat
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Mouse, Rat
Rabbit
Rabbit
Polyclonal
Rabbit IgG
250-277

DDX39 Antibody (Center Y265) - Additional Information

Gene ID 10212

Other Names

ATP-dependent RNA helicase DDX39A, DEAD box protein 39, Nuclear RNA helicase URH49, DDX39A, DDX39

Target/Specificity

This DDX39 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 250-277 amino acids from the Central region of human DDX39.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DDX39 Antibody (Center Y265) is for research use only and not for use in diagnostic or therapeutic procedures.

DDX39 Antibody (Center Y265) - Protein Information

Name DDX39A





Synonyms DDX39

Function [Isoform 1]: Involved in pre-mRNA splicing. Required for the export of mRNA out of the nucleus.

Cellular Location

Nucleus. Cytoplasm. Note=Can translocate to the cytoplasm in the presence of MX1

Tissue Location


Detected in testis, and at lower levels in brain, kidney, lung, thymus, spleen and salivary gland

DDX39 Antibody (Center Y265) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

DDX39 Antibody (Center Y265) - Images



Western blot analysis of DDX39 Antibody (Center Y265) (Cat. #AP7451d) in HepG2 cell line lysates (35ug/lane). DDX39 (arrow) was detected using the purified Pab.





Western blot analysis of DDX39 (arrow) using rabbit polyclonal DDX39 Antibody (Center Y265) (Cat. #AP7451d). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the DDX39 gene.

DDX39 Antibody (Center Y265) - Background

DDX39 is a member of the DEAD box protein family. These proteins are characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD) and are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of the DEAD box protein family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division.

DDX39 Antibody (Center Y265) - References

Pryor A., Tung L., Yang Z.Nucleic Acids Res. 32:1857-1865(2004) Lehner B., Semple J.I., Brown S.E.Genomics 83:153-167(2004)