

**SNX5 Rabbit mAb**  
**Catalog # AP74878****Specification****SNX5 Rabbit mAb - Product Information**

Application	WB, IP, ICC
Primary Accession	<a href="#">Q9Y5X3</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	46816

**SNX5 Rabbit mAb - Additional Information****Gene ID** 27131**Other Names**

SNX5

**Dilution**

WB~~1/500-1/1000

IP~~1/20

ICC~~N/A

**Format**

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

**Storage**

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

**SNX5 Rabbit mAb - Protein Information****Name** SNX5**Function**

Involved in several stages of intracellular trafficking. Interacts with membranes containing phosphatidylinositol 3-phosphate (PtdIns(3P)) or phosphatidylinositol 3,4-bisphosphate (PtdIns(3,4)P2) (PubMed:<a href="http://www.uniprot.org/citations/15561769" target="\_blank">15561769</a>). Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC) (Probable). Does not have in vitro vesicle-to-membrane remodeling activity (PubMed:<a href="http://www.uniprot.org/citations/23085988" target="\_blank">23085988</a>). Involved in retrograde transport of lysosomal enzyme receptor IGF2R (PubMed:<a href="http://www.uniprot.org/citations/17148574" target="\_blank">17148574</a>, PubMed:<a href="http://www.uniprot.org/citations/18596235" target="\_blank">18596235</a>). May function

as link between endosomal transport vesicles and dynactin (Probable). Plays a role in the internalization of EGFR after EGF stimulation (Probable). Involved in EGFR endosomal sorting and degradation; the function involves PIP5K1C isoform 3 and is retromer- independent (PubMed:<a href="http://www.uniprot.org/citations/23602387" target="\_blank">23602387</a>). Together with PIP5K1C isoform 3 facilitates HGS interaction with ubiquitinated EGFR, which initiates EGFR sorting to intraluminal vesicles (ILVs) of the multivesicular body for subsequent lysosomal degradation (Probable). Involved in E-cadherin sorting and degradation; inhibits PIP5K1C isoform 3-mediated E-cadherin degradation (PubMed:<a href="http://www.uniprot.org/citations/24610942" target="\_blank">24610942</a>). Plays a role in macropinocytosis (PubMed:<a href="http://www.uniprot.org/citations/18854019" target="\_blank">18854019</a>, PubMed:<a href="http://www.uniprot.org/citations/21048941" target="\_blank">21048941</a>).

### Cellular Location

Endosome. Early endosome Early endosome membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm. Cell projection, phagocytic cup. Cell projection, ruffle. Note=Recruited to the plasma membrane after EGF stimulation, which leads to increased levels of phosphatidylinositol 3,4-bisphosphate (PtdIns(3,4)P<sub>2</sub>) (PubMed:15561769). Detected on macropinosomes (PubMed:16968745, PubMed:21048941). Targeted to membrane ruffles in response to EGFR stimulation.

### SNX5 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### SNX5 Rabbit mAb - Images



