

**EIF5A Rabbit mAb**  
**Catalog # AP75011****Specification**

---

**EIF5A Rabbit mAb - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P, IP          |
| Primary Accession | <a href="#">P63241</a> |
| Reactivity        | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Monoclonal Antibody    |
| Calculated MW     | 16832                  |

**EIF5A Rabbit mAb - Additional Information****Gene ID** 1984**Other Names**  
EIF5A**Dilution**  
WB~~1/500-1/1000  
IHC-P~~N/A  
IP~~1/20**Format**  
Liquid**EIF5A Rabbit mAb - Protein Information****Name** EIF5A ([HGNC:3300](#))**Function**

Translation factor that promotes translation elongation and termination, particularly upon ribosome stalling at specific amino acid sequence contexts (PubMed:<a href="http://www.uniprot.org/citations/33547280" target="\_blank">33547280</a>). Binds between the exit (E) and peptidyl (P) site of the ribosome and promotes rescue of stalled ribosome: specifically required for efficient translation of polyproline-containing peptides as well as other motifs that stall the ribosome (By similarity). Acts as a ribosome quality control (RQC) cofactor by joining the RQC complex to facilitate peptidyl transfer during CAT tailing step (By similarity). Also involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall integrity (PubMed:<a href="http://www.uniprot.org/citations/16987817" target="\_blank">16987817</a>). With syntenin SDCBP, functions as a regulator of p53/TP53 and p53/TP53-dependent apoptosis (PubMed:<a href="http://www.uniprot.org/citations/15371445" target="\_blank">15371445</a>). Also regulates TNF-alpha-mediated apoptosis (PubMed:<a href="http://www.uniprot.org/citations/15452064" target="\_blank">15452064</a>, PubMed:<a href="http://www.uniprot.org/citations/17187778" target="\_blank">17187778</a>). Mediates effects of polyamines on neuronal process extension and survival (PubMed:<a

[17360499](http://www.uniprot.org/citations/17360499)). Is required for autophagy by assisting the ribosome in translating the ATG3 protein at a specific amino acid sequence, the 'ASP-ASP-Gly' motif, leading to the increase of the efficiency of ATG3 translation and facilitation of LC3B lipidation and autophagosome formation (PubMed:[29712776](http://www.uniprot.org/citations/29712776)).

#### Cellular Location

Cytoplasm. Nucleus. Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Note=Hypusine modification promotes the nuclear export and cytoplasmic localization and there was a dynamic shift in the localization from predominantly cytoplasmic to primarily nuclear under apoptotic inducing conditions (PubMed:19379712, PubMed:27306458). Nuclear export of hypusinated protein is mediated by XPO4 (PubMed:10944119, PubMed:27306458).

#### Tissue Location

Expressed in umbilical vein endothelial cells and several cancer cell lines (at protein level)

### eIF5A Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### eIF5A Rabbit mAb - Images



