

67kDa Laminin Receptor Rabbit mAb

Catalog # AP75019

Specification

67kDa Laminin Receptor Rabbit mAb - Product Information

Application WB, IP Primary Accession P08865

Reactivity Human, Mouse, Rat

Host Rabb

Clonality Monoclonal Antibody

Calculated MW 32854

67kDa Laminin Receptor Rabbit mAb - Additional Information

Gene ID 3921

Other Names

RPSA

Dilution

WB~~1/500-1/1000

 $IP \sim \sim N/A$

Format

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

67kDa Laminin Receptor Rabbit mAb - Protein Information

Name RPSA {ECO:0000255|HAMAP-Rule:MF_03016}

Synonyms LAMBR, LAMR1

Function

Required for the assembly and/or stability of the 40S ribosomal subunit. Required for the processing of the 20S rRNA- precursor to mature 18S rRNA in a late step of the maturation of 40S ribosomal subunits. Also functions as a cell surface receptor for laminin. Plays a role in cell adhesion to the basement membrane and in the consequent activation of signaling transduction pathways. May play a role in cell fate determination and tissue morphogenesis. Acts as a PPP1R16B-dependent substrate of PPP1CA.

Cellular Location

Cell membrane. Cytoplasm. Nucleus {ECO:0000255|HAMAP-Rule:MF_03016}. Note=67LR is found at the surface of the plasma membrane, with its C-terminal laminin-binding domain accessible to extracellular ligands. 37LRP is found at the cell surface, in the cytoplasm and in the nucleus (By similarity) Colocalizes with PPP1R16B in the cell membrane. {ECO:0000255|HAMAP-



Rule:MF_03016}

67kDa Laminin Receptor Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

67kDa Laminin Receptor Rabbit mAb - Images

