

ApoER2 Rabbit mAb

Catalog # AP75092

Specification

ApoER2 Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, IP
O14114
Human, Rat, Hamster
Rabbit
Monoclonal Antibody
105634

ApoER2 Rabbit mAb - Additional Information

Gene ID 7804

Other Names LRP8

DilutionWB~~1/500-1/1000
IP~~1/20

Format

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

ApoER2 Rabbit mAb - Protein Information

Name LRP8

Synonyms APOER2

Function

Cell surface receptor for Reelin (RELN) and apolipoprotein E (apoE)-containing ligands (PubMed:12899622, PubMed:12899622, PubMed:20223215, PubMed:30873003). LRP8 participates in transmitting the extracellular Reelin signal to intracellular signaling processes, by binding to DAB1 on its cytoplasmic tail (By similarity). Reelin acts via both the VLDL receptor (VLDLR) and LRP8 to regulate DAB1 tyrosine phosphorylation and microtubule function in neurons (By similarity). LRP8 has higher affinity for Reelin than VLDLR (By similarity). LRP8 is thus a key component of the Reelin pathway which governs neuronal layering of the forebrain during embryonic brain development (By similarity). Binds the endoplasmic reticulum resident receptor-associated protein (RAP) (By similarity). Binds dimers of beta 2- glycoprotein I and may be



involved in the suppression of platelet aggregation in the vasculature (PubMed:12807892). Highly expressed in the initial segment of the epididymis, where it affects the functional expression of clusterin and phospholipid hydroperoxide glutathione peroxidase (PHGPx), two proteins required for sperm maturation (By similarity). May also function as an endocytic receptor (By similarity). Not required for endocytic uptake of SEPP1 in the kidney which is mediated by LRP2 (By similarity). Together with its ligand, apolipoprotein E (apoE), may indirectly play a role in the suppression of the innate immune response by controlling the survival of myeloid- derived suppressor cells (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Secreted {ECO:0000250|UniProtKB:Q924X6}. Note=Isoforms that contain the exon coding for a furin-type cleavage site are proteolytically processed, leading to a secreted receptor fragment {ECO:0000250|UniProtKB:Q924X6}

Tissue Location

Expressed mainly in brain and placenta. Also expressed in platelets and megakaryocytic cells. Not expressed in the liver.

ApoER2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ApoER2 Rabbit mAb - Images





