

**BMP2 Rabbit mAb**  
**Catalog # AP75160****Specification**

---

**BMP2 Rabbit mAb - Product Information**

Application	WB, ICC
Primary Accession	<a href="#">P12643</a>
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	44702

**BMP2 Rabbit mAb - Additional Information****Gene ID** 650**Other Names**  
BMP2**Dilution**  
WB~~1/500-1/1000  
ICC~~N/A**Format**  
Liquid**BMP2 Rabbit mAb - Protein Information****Name** BMP2**Synonyms** BMP2A**Function**

Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes, including cardiogenesis, neurogenesis, and osteogenesis (PubMed:<a href="http://www.uniprot.org/citations/18436533" target="\_blank">18436533</a>, PubMed:<a href="http://www.uniprot.org/citations/24362451" target="\_blank">24362451</a>, PubMed:<a href="http://www.uniprot.org/citations/31019025" target="\_blank">31019025</a>). Induces cartilage and bone formation (PubMed:<a href="http://www.uniprot.org/citations/3201241" target="\_blank">3201241</a>). Initiates the canonical BMP signaling cascade by associating with type I receptor BMPRI1A and type II receptor BMPRI2 (PubMed:<a href="http://www.uniprot.org/citations/15064755" target="\_blank">15064755</a>, PubMed:<a href="http://www.uniprot.org/citations/17295905" target="\_blank">17295905</a>, PubMed:<a href="http://www.uniprot.org/citations/18436533" target="\_blank">18436533</a>). Once all three components are bound together in a complex at the cell surface, BMPRI2 phosphorylates and activates BMPRI1A (PubMed:<a href="http://www.uniprot.org/citations/7791754" target="\_blank">7791754</a>). In turn, BMPRI1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes. Also acts to promote expression of HAMP, via the interaction with its receptor

BMPR1A/ALK3 (PubMed:<a href="http://www.uniprot.org/citations/31800957" target="\_blank">31800957</a>). Can also signal through non-canonical pathways such as ERK/MAP kinase signaling cascade that regulates osteoblast differentiation (PubMed:<a href="http://www.uniprot.org/citations/16771708" target="\_blank">16771708</a>, PubMed:<a href="http://www.uniprot.org/citations/20851880" target="\_blank">20851880</a>). Also stimulates the differentiation of myoblasts into osteoblasts via the EIF2AK3-EIF2A-ATF4 pathway by stimulating EIF2A phosphorylation which leads to increased expression of ATF4 which plays a central role in osteoblast differentiation (PubMed:<a href="http://www.uniprot.org/citations/24362451" target="\_blank">24362451</a>). Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, expression is repressed during the bell stage by MSX1-mediated inhibition of CTNNB1 signaling (By similarity).

#### Cellular Location

Secreted.

#### Tissue Location

Particularly abundant in lung, spleen and colon and in low but significant levels in heart, brain, placenta, liver, skeletal muscle, kidney, pancreas, prostate, ovary and small intestine

#### BMP2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### BMP2 Rabbit mAb - Images

