

**CD3D Rabbit mAb**  
**Catalog # AP75228****Specification****CD3D Rabbit mAb - Product Information**

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	<a href="#">P04234</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	18930

**CD3D Rabbit mAb - Additional Information****Gene ID 915****Other Names**  
CD3D**Dilution**

WB~~1/500-1/1000  
IHC-P~~N/A  
IHC-F~~N/A  
IP~~N/A  
ICC~~N/A

**Format**

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

**Storage**

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

**CD3D Rabbit mAb - Protein Information****Name** CD3D**Synonyms** T3D**Function**

Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR- mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:[2470098](http://www.uniprot.org/citations/2470098)). In addition of this role of signal transduction in T-cell activation, CD3D plays an essential role in thymocyte differentiation. Indeed, participates in correct

intracellular TCR-CD3 complex assembly and surface expression. In absence of a functional TCR-CD3 complex, thymocytes are unable to differentiate properly. Interacts with CD4 and CD8 and thus serves to establish a functional link between the TCR and coreceptors CD4 and CD8, which is needed for activation and positive selection of CD4 or CD8 T-cells (PubMed:<a href="http://www.uniprot.org/citations/12215456" target="\_blank">12215456</a>).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein

#### Tissue Location

CD3D is mostly present on T-lymphocytes with its TCR-CD3 partners. Present also in fetal NK-cells

#### CD3D Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### CD3D Rabbit mAb - Images

