

CENPE Rabbit mAb
Catalog # AP75255**Specification**

CENPE Rabbit mAb - Product Information

Application	WB
Primary Accession	Q02224
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	316415

CENPE Rabbit mAb - Additional Information**Gene ID** 1062**Other Names**
CENPE**Dilution**
WB~~1/500-1/1000**Format**
Liquid**CENPE Rabbit mAb - Protein Information****Name** CENPE**Function**

Microtubule plus-end-directed kinetochore motor which plays an important role in chromosome congression, microtubule-kinetochore conjugation and spindle assembly checkpoint activation. Drives chromosome congression (alignment of chromosomes at the spindle equator resulting in the formation of the metaphase plate) by mediating the lateral sliding of polar chromosomes along spindle microtubules towards the spindle equator and by aiding the establishment and maintenance of connections between kinetochores and spindle microtubules (PubMed:23891108, PubMed:25395579, PubMed:7889940). The transport of pole-proximal chromosomes towards the spindle equator is favored by microtubule tracks that are deetyrosinated (PubMed:25908662). Acts as a processive bi-directional tracker of dynamic microtubule tips; after chromosomes have congressed, continues to play an active role at kinetochores, enhancing their links with dynamic microtubule ends (PubMed:23955301). Suppresses chromosome congression in NDC80-depleted cells and contributes positively to congression only when microtubules are stabilized (PubMed:25743205). Plays an important role in the formation of stable attachments

between kinetochores and spindle microtubules (PubMed:17535814) The stabilization of kinetochore- microtubule attachment also requires CENPE-dependent localization of other proteins to the kinetochore including BUB1B, MAD1 and MAD2. Plays a role in spindle assembly checkpoint activation (SAC) via its interaction with BUB1B resulting in the activation of its kinase activity, which is important for activating SAC. Necessary for the mitotic checkpoint signal at individual kinetochores to prevent aneuploidy due to single chromosome loss (By similarity).

Cellular Location

Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, spindle. Chromosome, centromere. Note=Associates with kinetochores during congression (as early as prometaphase), relocates to the spindle midzone at anaphase, and is quantitatively discarded at the end of the cell division (By similarity). Recruited to the kinetochore in a SEPT7, CENPQ and TRAPPC12-dependent manner (PubMed:18460473, PubMed:25395579, PubMed:25918224). Recruited to the pericentromeric/centromeric regions of the chromosome in a CTCF- dependent manner (PubMed:26321640). {ECO:0000250|UniProtKB:Q6RT24, ECO:0000269|PubMed:18460473, ECO:0000269|PubMed:25395579, ECO:0000269|PubMed:25918224, ECO:0000269|PubMed:26321640}

CENPE Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CENPE Rabbit mAb - Images

