

Clusterin alpha Chain Rabbit mAb
Catalog # AP75274**Specification****Clusterin alpha Chain Rabbit mAb - Product Information**

Application	WB, IP
Primary Accession	Q06890
Reactivity	Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	51656

Clusterin alpha Chain Rabbit mAb - Additional Information**Gene ID** 12759**Other Names**

Clu

Dilution

WB~~1/500-1/1000

IP~~N/A

Format

Liquid

Clusterin alpha Chain Rabbit mAb - Protein Information**Name** Clu {ECO:0000312|MGI:MGI:88423}**Function**

Functions as extracellular chaperone that prevents aggregation of non native proteins. Prevents stress-induced aggregation of blood plasma proteins (By similarity). Inhibits formation of amyloid fibrils by APP, APOC2, B2M, CALCA, CSN3, SNCA and aggregation-prone LYZ variants (in vitro) (PubMed:14741101). Does not require ATP. Maintains partially unfolded proteins in a state appropriate for subsequent refolding by other chaperones, such as HSPA8/HSC70. Does not refold proteins by itself. Binding to cell surface receptors triggers internalization of the chaperone-client complex and subsequent lysosomal or proteasomal degradation. When secreted, protects cells against apoptosis and against cytolysis by complement: inhibits assembly of the complement membrane attack complex (MAC) by preventing polymerization of C9 pore component of the MAC complex. Intracellular forms interact with ubiquitin and SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complexes and promote the ubiquitination and subsequent proteasomal degradation of target proteins. Promotes proteasomal degradation of COMMD1 and IKBKB. Modulates NF-kappa-B transcriptional activity (By similarity). Following stress, promotes apoptosis (PubMed:12551933). Inhibits apoptosis when associated with the mitochondrial membrane by interference with BAX-dependent release of cytochrome c into the cytoplasm. Plays a role in the regulation of cell proliferation.

Following ER stress, suppresses stress-induced apoptosis by stabilizing mitochondrial membrane integrity through interaction with HSPA5. When secreted, does not affect caspase or BAX-mediated intrinsic apoptosis and TNF-induced NF-kappa-B-activity (By similarity). When secreted, acts as an important modulator during neuronal differentiation through interaction with STMN3 (By similarity). Plays a role in the clearance of immune complexes that arise during cell injury (PubMed:11865066).

Cellular Location

Secreted. Nucleus. Cytoplasm Mitochondrion membrane {ECO:0000250|UniProtKB:P10909}; Peripheral membrane protein {ECO:0000250|UniProtKB:P10909}; Cytoplasmic side {ECO:0000250|UniProtKB:P10909}. Cytoplasm, cytosol. Microsome {ECO:0000250|UniProtKB:P10909}. Endoplasmic reticulum {ECO:0000250|UniProtKB:P10909}. Mitochondrion {ECO:0000250|UniProtKB:P10909}. Mitochondrion membrane {ECO:0000250|UniProtKB:P10909}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:P05371}. Cytoplasmic vesicle, secretory vesicle, chromaffin granule {ECO:0000250|UniProtKB:P10909}. Note=Can retrotranslocate from the secretory compartments to the cytosol upon cellular stress. Detected in perinuclear foci that may be aggresomes containing misfolded, ubiquitinated proteins. Detected at the mitochondrion membrane upon induction of apoptosis. Under ER stress, a immaturely glycosylated pre-secreted form retrotranslocates from the endoplasmic reticulum (ER)-Golgi network to the cytoplasm to localize in the mitochondria through HSPA5 interaction. ER stress reduces secretion. Under the stress, minor amounts of non-secreted forms accumulate in cytoplasm. {ECO:0000250|UniProtKB:P10909}

Tissue Location

Most abundant in stomach, liver, brain, and testis, with intermediate levels in heart, ovary and kidney

Clusterin alpha Chain Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Clusterin alpha Chain Rabbit mAb - Images



