

Cyclin E2 Rabbit mAb

Catalog # AP75315

Specification

Cyclin E2 Rabbit mAb - Product Information

Application WB, IP, ICC
Primary Accession
Reactivity Human
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 46757

Cyclin E2 Rabbit mAb - Additional Information

Gene ID 9134

Other Names

CCNE2

Dilution

WB~~1/500-1/1000

IP~~N/A ICC~~N/A

Format

Liquid

Cyclin E2 Rabbit mAb - Protein Information

Name CCNE2

Function

Essential for the control of the cell cycle at the late G1 and early S phase.

Cellular Location

Nucleus.

Tissue Location

According to PubMed:9858585, highest levels of expression in adult testis, thymus and brain. Lower levels in placenta, spleen and colon. Consistently elevated levels in tumor-derived cells compared to non-transformed proliferating cells. According to PubMed:9840927: low levels in thymus, prostate, brain, skeletal muscle, and kidney. Elevated levels in lung. According to PubMed:9840943 highly expressed in testis, placenta, thymus and brain. In a lesser extent in small intestine and colon.

Cyclin E2 Rabbit mAb - Protocols



Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Cyclin E2 Rabbit mAb - Images



