

DcR2 Rabbit mAb
Catalog # AP75345**Specification**

DcR2 Rabbit mAb - Product Information

Application	WB, IHC-P
Primary Accession	Q9UBN6
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	41849

DcR2 Rabbit mAb - Additional Information**Gene ID** 8793**Other Names**
TNFRSF10D**Dilution**
WB~~1/500-1/1000
IHC-P~~N/A**Format**
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**DcR2 Rabbit mAb - Protein Information****Name** TNFRSF10D ([HGNC:11907](#))**Function**

Receptor for the cytotoxic ligand TRAIL (PubMed:9430226). Contains a truncated death domain and hence is not capable of inducing apoptosis but protects against TRAIL-mediated apoptosis (PubMed:9537512). Reports are contradictory with regards to its ability to induce the NF-kappa-B pathway. According to PubMed:9382840, it cannot but according to PubMed:9430226, it can induce the NF-kappa-B pathway (PubMed:9382840, PubMed:9430226).

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location

Widely expressed, in particular in fetal kidney, lung and liver, and in adult testis and liver. Also expressed in peripheral blood leukocytes, colon and small intestine, ovary, prostate, thymus, spleen, pancreas, kidney, lung, placenta and heart

DcR2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DcR2 Rabbit mAb - Images

