

HMGCL Rabbit mAb
Catalog # AP75555**Specification**

HMGCL Rabbit mAb - Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | P35914 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Monoclonal Antibody |
| Calculated MW | 34360 |

HMGCL Rabbit mAb - Additional Information**Gene ID** 3155**Other Names**
HMGCL**Dilution**
WB~~1/500-1/1000**Format**
Liquid**HMGCL Rabbit mAb - Protein Information****Name** HMGCL**Function**

Mitochondrial 3-hydroxy-3-methylglutaryl-CoA lyase that catalyzes a cation-dependent cleavage of (S)-3-hydroxy-3-methylglutaryl-CoA into acetyl-CoA and acetoacetate, a key step in ketogenesis. Terminal step in leucine catabolism. Ketone bodies (beta-hydroxybutyrate, acetoacetate and acetone) are essential as an alternative source of energy to glucose, as lipid precursors and as regulators of metabolism.

Cellular Location

Mitochondrion matrix {ECO:0000250|UniProtKB:P38060}. Peroxisome {ECO:0000250|UniProtKB:P38060}. Note=Unprocessed form is peroxisomal {ECO:0000250|UniProtKB:P38060}

Tissue Location

Highest expression in liver. Expressed in pancreas, kidney, intestine, testis, fibroblasts and lymphoblasts. Very low expression in brain and skeletal muscle. The relative expression of isoform 2 (at mRNA level) is highest in heart (30%), skeletal muscle (22%), and brain (14%).

HMGCL Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HMGCL Rabbit mAb - Images

