

Hsp27 Rabbit mAb

Catalog # AP75580

Specification

Hsp27 Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW

WB, IHC-P, IHC-F, IP, ICC P04792 Human, Mouse Rabbit **Monoclonal Antibody** 22783

Hsp27 Rabbit mAb - Additional Information

Gene ID 3315

Other Names HSPB1

Dilution

WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~N/A ICC~~N/A

Format Liquid

Hsp27 Rabbit mAb - Protein Information

Name HSPB1

Synonyms HSP27, HSP28

Function

Small heat shock protein which functions as a molecular chaperone probably maintaining denatured proteins in a folding- competent state (PubMed: <a

 $href="http://www.uniprot.org/citations/10383393" target="_blank">10383393, PubMed:20178975). Plays a role$ in stress resistance and actin organization (PubMed:19166925). Through its molecular chaperone activity may regulate numerous biological processes including the phosphorylation and the axonal transport of neurofilament proteins (PubMed:23728742).

Cellular Location

Cytoplasm. Nucleus Cytoplasm, cytoskeleton, spindle Note=Cytoplasmic in interphase cells.





Tel: 858.875.1900 Fax: 858.875.1999

Colocalizes with mitotic spindles in mitotic cells. Translocates to the nucleus during heat shock and resides in sub-nuclear structures known as SC35 speckles or nuclear splicing speckles.

Tissue Location

Detected in all tissues tested: skeletal muscle, heart, aorta, large intestine, small intestine, stomach, esophagus, bladder, adrenal gland, thyroid, pancreas, testis, adipose tissue, kidney, liver, spleen, cerebral cortex, blood serum and cerebrospinal fluid. Highest levels are found in the heart and in tissues composed of striated and smooth muscle.

Hsp27 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Hsp27 Rabbit mAb - Images





