

ICOS Ligand Rabbit mAb

Catalog # AP75591

Specification

ICOS Ligand Rabbit mAb - Product Information

Application WB, IP
Primary Accession O75144
Reactivity Human
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 33349

ICOS Ligand Rabbit mAb - Additional Information

Gene ID 23308

Other Names ICOSLG

DilutionWB~~1/500-1/1000
IP~~N/A

Format Liquid

ICOS Ligand Rabbit mAb - Protein Information

Name ICOSLG

Function

Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion (PubMed:11007762, PubMed:11023515, PubMed:30498080). Also induces B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function (By similarity). In endothelial cells, required for proper neutrophil transmigration in response to chemoattractants, such as CXCL8/IL8 or N-formyl-methionyl peptides (fMLP) (PubMed:30498080).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed on peripheral blood B-cells and monocytes, as well as on monocyte-derived dendritic cells (at protein level). [Isoform 2]: Detected only in lymph nodes, leukocytes and spleen.



Expressed on activated monocytes and dendritic cells.

ICOS Ligand Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ICOS Ligand Rabbit mAb - Images

