

IRF8 Rabbit mAb

Catalog # AP75632

Specification

IRF8 Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC-P, IP <u>002556</u> Human Rabbit Monoclonal Antibody 48356

IRF8 Rabbit mAb - Additional Information

Gene ID 3394

Other Names IRF8

Dilution WB~~1/500-1/1000 IHC-P~~N/A IP~~N/A

Format Liquid

IRF8 Rabbit mAb - Protein Information

Name IRF8 {ECO:0000303|PubMed:21524210, ECO:0000312|HGNC:HGNC:5358}

Function

Transcription factor that specifically binds to the upstream regulatory region of type I interferon (IFN) and IFN-inducible MHC class I genes (the interferon consensus sequence (ICS)) (PubMed:25122610). Can both act as a transcriptional activator or repressor (By similarity). Plays a negative regulatory role in cells of the immune system (By similarity). Involved in CD8(+) dendritic cell differentiation by forming a complex with the BATF-JUNB heterodimer in immune cells, leading to recognition of AICE sequence (5'-TGAnTCA/GAAA- 3'), an immune-specific regulatory element, followed by cooperative binding of BATF and IRF8 and activation of genes (By similarity). Required for the development of plasmacytoid dendritic cells (pDCs), which produce most of the type I IFN in response to viral infection (By similarity). Positively regulates macroautophagy in dendritic cells (PubMed:29434592). Acts as a transcriptional repressor of osteoclast differentiation factors such as NFATC1 and EEIG1 (By similarity).

Cellular Location

Nucleus. Cytoplasm Note=In resting macrophages, localizes in the cytoplasm. Translocated in the



nucleus upon IFN-gamma induction.

Tissue Location Predominantly expressed in lymphoid tissues.

IRF8 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

IRF8 Rabbit mAb - Images

