

CLIC4 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7564a

Specification

CLIC4 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Isotype Calculated MW IHC-P, WB,E <u>O9Y696</u> Human Rabbit Polyclonal Rabbit IgG 28772

CLIC4 Antibody - Additional Information

Gene ID 25932

Other Names Chloride intracellular channel protein 4, Intracellular chloride ion channel protein p64H1, CLIC4

Target/Specificity This CLIC4 antibody is generated from rabbits immunized with recombinant human CLIC4 protein.

Dilution IHC-P~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions CLIC4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CLIC4 Antibody - Protein Information

Name CLIC4 {ECO:0000303|PubMed:12163372, ECO:0000312|HGNC:HGNC:13518}

Function In the soluble state, catalyzes glutaredoxin-like thiol disulfide exchange reactions with reduced glutathione as electron donor (PubMed:<u>25581026</u>, PubMed:<u>37759794</u>). Can insert into membranes and form voltage-dependent multi-ion conductive channels. Membrane insertion seems to be redox-regulated and may occur only under oxidizing conditions (By similarity)



(PubMed:<u>16176272</u>). Has alternate cellular functions like a potential role in angiogenesis or in maintaining apical-basolateral membrane polarity during mitosis and cytokinesis. Could also promote endothelial cell proliferation and regulate endothelial morphogenesis (tubulogenesis). Promotes cell-surface expression of HRH3.

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasmic vesicle membrane; Single-pass membrane protein. Nucleus. Cell membrane; Single-pass membrane protein. Mitochondrion {ECO:0000250|UniProtKB:Q9Z0W7}. Cell junction. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q9Z0W7}; Single-pass membrane protein {ECO:0000250|UniProtKB:Q9Z0W7}. Note=Colocalized with AKAP9 at the centrosome and midbody. Exists both as soluble cytoplasmic protein and as membrane protein with probably a single transmembrane domain Present in an intracellular vesicular compartment that likely represent trans-Golgi network vesicles. Might not be present in the nucleus of cardiac cells. {ECO:0000250|UniProtKB:Q9Z0W7, ECO:0000269|PubMed:14569596}

Tissue Location

Detected in epithelial cells from colon, esophagus and kidney (at protein level). Expression is prominent in heart, kidney, placenta and skeletal muscle.

CLIC4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

CLIC4 Antibody - Images



Western blot analysis of CLIC4 (arrow) using rabbit polyclonal CLIC4 Antibody (Cat.#AP7564a).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the CLIC4 gene (Lane 2) (Origene Technologies).





Formalin-fixed and paraffin-embedded human breast carcinoma tissue reacted with CLIC4 antibody (Cat.#AP7564a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

CLIC4 Antibody - Background

Chloride channels are a diverse group of proteins that regulate fundamental cellular processes including stabilization of cell membrane potential, transpithelial transport, maintenance of intracellular pH, and regulation of cell volume. Chloride intracellular channel 4 (CLIC4) protein, encoded by the CLIC4 gene, is a member of the p64 family; the gene is expressed in many tissues and exhibits a intracellular vesicular pattern in Panc-1 cells (pancreatic cancer cells).

CLIC4 Antibody - References

Singh,H.,FEBS J. 274 (24), 6306-6316 (2007) Suh,K.S.,J. Cell. Sci. 120 (PT 15), 2631-2640 (2007) Suh,K.S.,Clin. Cancer Res. 13 (1), 121-131 (2007) CLIC4 Antibody - Citations

• Association of chloride intracellular channel 4 and Indian hedgehog proteins with survival of patients with pancreatic ductal adenocarcinoma.