

Phospho-LAT (Tyr200) Rabbit mAb
Catalog # AP75667**Specification**

Phospho-LAT (Tyr200) Rabbit mAb - Product Information

Application	WB
Primary Accession	O43561
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	27930

Phospho-LAT (Tyr200) Rabbit mAb - Additional Information**Gene ID** 27040**Other Names**

LAT

Dilution

WB~~1/500-1/1000

Format

Liquid

Phospho-LAT (Tyr200) Rabbit mAb - Protein Information**Name** LAT**Function**

Required for TCR (T-cell antigen receptor)- and pre-TCR- mediated signaling, both in mature T-cells and during their development (PubMed:<[a href="http://www.uniprot.org/citations/23514740" target="_blank">23514740](http://www.uniprot.org/citations/23514740), PubMed:<[a href="http://www.uniprot.org/citations/25907557" target="_blank">25907557](http://www.uniprot.org/citations/25907557)). Involved in FCGR3 (low affinity immunoglobulin gamma Fc region receptor III)-mediated signaling in natural killer cells and FCER1 (high affinity immunoglobulin epsilon receptor)-mediated signaling in mast cells. Couples activation of these receptors and their associated kinases with distal intracellular events such as mobilization of intracellular calcium stores, PKC activation, MAPK activation or cytoskeletal reorganization through the recruitment of PLCG1, GRB2, GRAP2, and other signaling molecules.

Cellular Location

Cell membrane; Single-pass type III membrane protein. Note=Present in lipid rafts

Tissue Location

Expressed in thymus, T-cells, NK cells, mast cells and, at lower levels, in spleen. Present in T-cells but not B-cells (at protein level).

Phospho-LAT (Tyr200) Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-LAT (Tyr200) Rabbit mAb - Images

