

**Phospho-MCM2 (Ser108) Rabbit mAb**  
**Catalog # AP75696****Specification****Phospho-MCM2 (Ser108) Rabbit mAb - Product Information**

Application	WB, IP, ICC
Primary Accession	<a href="#">P49736</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	101896

**Phospho-MCM2 (Ser108) Rabbit mAb - Additional Information****Gene ID** 4171**Other Names**  
MCM2**Dilution**  
WB~~1/500-1/1000  
IP~~1/20  
ICC~~N/A**Format**  
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**  
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**Phospho-MCM2 (Ser108) Rabbit mAb - Protein Information****Name** MCM2 ([HGNC:6944](#))**Function**

Acts as a component of the MCM2-7 complex (MCM complex) which is the replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Core component of CDC45-MCM-GINS (CMG) helicase, the molecular machine that unwinds template DNA during replication, and around which the replisome is built (PubMed:<a href="http://www.uniprot.org/citations/32453425" target="\_blank">32453425</a>, PubMed:<a href="http://www.uniprot.org/citations/34694004" target="\_blank">34694004</a>, PubMed:<a href="http://www.uniprot.org/citations/34700328" target="\_blank">34700328</a>, PubMed:<a href="http://www.uniprot.org/citations/35585232" target="\_blank">35585232</a>). The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity (PubMed:<a href="http://www.uniprot.org/citations/32453425" target="\_blank">32453425</a>). Required for

the entry in S phase and for cell division (PubMed:<a href="http://www.uniprot.org/citations/8175912" target="\_blank">8175912</a>). Plays a role in terminally differentiated hair cells development of the cochlea and induces cells apoptosis (PubMed:<a href="http://www.uniprot.org/citations/26196677" target="\_blank">26196677</a>).

#### Cellular Location

Nucleus. Chromosome. Note=Associated with chromatin before the formation of nuclei and detaches from it as DNA replication progresses. {ECO:0000250|UniProtKB:P55861}

#### Phospho-MCM2 (Ser108) Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Phospho-MCM2 (Ser108) Rabbit mAb - Images



