

NCE2 Rabbit mAb
Catalog # AP75772**Specification**

NCE2 Rabbit mAb - Product Information

Application	WB, IHC-P, IHC-F, ICC
Primary Accession	Q969M7
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	21077

NCE2 Rabbit mAb - Additional Information**Gene ID** 140739**Other Names**
UBE2F**Dilution**
WB~~1/500-1/1000
IHC-P~~N/A
IHC-F~~N/A
ICC~~N/A**Format**
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**NCE2 Rabbit mAb - Protein Information****Name** UBE2F**Synonyms** NCE2**Function**

Accepts the ubiquitin-like protein NEDD8 from the UBA3-NAE1 E1 complex and catalyzes its covalent attachment to other proteins (PubMed:19250909, PubMed:23201271). Together with the E3 ubiquitin ligase RNF7/RBX2, specifically neddylates cullin-5 (CUL5) (PubMed:19250909, PubMed:23201271, PubMed:23300442). Does not neddylate CUL1, CUL2, CUL3, CUL4A or CUL4B (PubMed:19250909, PubMed:19250909, PubMed:19250909).

href="http://www.uniprot.org/citations/23201271" target="_blank">23201271). Mediates neddylation of the CUL9-RBX1 complex (PubMed:38605244).

Tissue Location

Widely expressed (at protein level).

NCE2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCE2 Rabbit mAb - Images