

NEK7 Rabbit mAb
Catalog # AP75793**Specification****NEK7 Rabbit mAb - Product Information**

Application	WB, IP
Primary Accession	Q8TDX7
Reactivity	Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	34551

NEK7 Rabbit mAb - Additional Information**Gene ID** 140609**Other Names**
NEK7**Dilution**
WB~~1/500-1/1000
IP~~1/20**Format**
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**NEK7 Rabbit mAb - Protein Information****Name** NEK7 {ECO:0000303|PubMed:11701951, ECO:0000312|HGNC:HGNC:13386}**Function**

Protein kinase which plays an important role in mitotic cell cycle progression (PubMed:17101132, PubMed:19941817, PubMed:31409757). Required for microtubule nucleation activity of the centrosome, robust mitotic spindle formation and cytokinesis (PubMed:17586473, PubMed:19414596, PubMed:19941817, PubMed:26522158, PubMed:31409757). Phosphorylates EML4 at 'Ser-146', promoting its dissociation from microtubules during mitosis which is required for efficient chromosome congression (PubMed:31409757). Phosphorylates RPS6KB1 (By similarity). Acts as an essential activator of the NLRP3 inflammasome

assembly independently of its kinase activity (PubMed:26642356, PubMed:36442502, PubMed:39173637). Acts by unlocking NLRP3 following NLRP3 translocation into the microtubule organizing center (MTOC), relieving NLRP3 autoinhibition and promoting formation of the NLRP3:PYCARD complex, and activation of CASP1 (PubMed:26642356, PubMed:31189953, PubMed:36442502, PubMed:39173637). Serves as a cellular switch that enforces mutual exclusivity of the inflammasome response and cell division: interaction with NEK9 prevents interaction with NLRP3 and activation of the inflammasome during mitosis (PubMed:26642356, PubMed:31189953).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9ES74}. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Present at centrosome throughout the cell cycle (PubMed:17586473). Also detected at spindle midzone of the anaphase cells and eventually concentrates at the midbody (PubMed:17586473). Interaction with ANKS3 prevents its translocation to the nucleus (By similarity). {ECO:0000250|UniProtKB:Q9ES74, ECO:0000269|PubMed:17586473}

Tissue Location

Highly expressed in lung, muscle, testis, brain, heart, liver, leukocyte and spleen. Lower expression in ovary, prostate and kidney. No expression seen in small intestine

NEK7 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NEK7 Rabbit mAb - Images



