

### **NOLC1** Rabbit mAb

**Catalog # AP75816** 

## **Specification**

#### **NOLC1** Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, IHC-P, IHC-F, ICC <u>Q14978</u> Human Rabbit Monoclonal Antibody 73603

#### NOLC1 Rabbit mAb - Additional Information

**Gene ID 9221** 

Other Names NOLC1

Dilution
WB~~1/500-1/1000
IHC-P~~N/A
IHC-F~~N/A
ICC~~N/A

Format Liquid

### **NOLC1 Rabbit mAb - Protein Information**

Name NOLC1 (HGNC:15608)

### **Function**

Nucleolar protein that acts as a regulator of RNA polymerase I by connecting RNA polymerase I with enzymes responsible for ribosomal processing and modification (PubMed:<a href="http://www.uniprot.org/citations/10567578" target="\_blank">10567578</a>, PubMed:<a href="http://www.uniprot.org/citations/26399832" target="\_blank">26399832</a>). Required for neural crest specification: following monoubiquitination by the BCR(KBTBD8) complex, associates with TCOF1 and acts as a platform to connect RNA polymerase I with enzymes responsible for ribosomal processing and modification, leading to remodel the translational program of differentiating cells in favor of neural crest specification (PubMed:<a href="http://www.uniprot.org/citations/26399832" target="\_blank">26399832</a>). Involved in nucleologenesis, possibly by playing a role in the maintenance of the fundamental structure of the fibrillar center and dense fibrillar component in the nucleolus (PubMed:<a href="http://www.uniprot.org/citations/9016786" target="\_blank">9016786</a>). It has intrinsic GTPase and ATPase activities (PubMed:<a href="http://www.uniprot.org/citations/9016786" target="\_blank">9016786</a>).





Tel: 858.875.1900 Fax: 858.875.1999

# **Cellular Location**

Nucleus, nucleolus. Cytoplasm. Note=Shuttles between the nucleolus and the cytoplasm. At telophase it begins to assemble into granular-like pre-nucleolar bodies which are subsequently relocated to nucleoli at the early G1-phase.

## **NOLC1** Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# NOLC1 Rabbit mAb - Images







