

# NUR77 Rabbit mAb

Catalog # AP75835

## Specification

## NUR77 Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB <u>P22736</u> Rat Rabbit Monoclonal Antibody 64463

## NUR77 Rabbit mAb - Additional Information

Gene ID 3164

**Other Names** NR4A1

**Dilution** WB~~1/500-1/1000

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

## Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## NUR77 Rabbit mAb - Protein Information

Name NR4A1

Synonyms GFRP1, HMR, NAK1

#### Function

Orphan nuclear receptor. Binds the NGFI-B response element (NBRE) 5'-AAAGGTCA-3' (PubMed:<a href="http://www.uniprot.org/citations/18690216" target="\_blank">18690216</a>, PubMed:<a href="http://www.uniprot.org/citations/8121493" target="\_blank">8121493</a>, PubMed:<a href="http://www.uniprot.org/citations/9315652" target="\_blank">9315652</a>). Binds 9-cis-retinoic acid outside of its ligand- binding (NR LBD) domain (PubMed:<a href="http://www.uniprot.org/citations/18690216" target="\_blank">18690216</a>). Participates in energy homeostasis by sequestrating the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation (PubMed:<a href="http://www.uniprot.org/citations/22983157" target="\_blank">22983157</a>). Regulates the inflammatory response in macrophages by regulating metabolic adaptations during inflammation, including repressing the transcription of genes involved in the citric acid cycle (TCA) (By similarity). Inhibits NF-kappa-B signaling by binding to low-affinity NF-kappa-B binding sites, such as at the IL2 promoter (PubMed:<a href="http://www.uniprot.org/citations/15466594" target="\_blank">15466594</a>). May act



concomitantly with NR4A2 in regulating the expression of delayed-early genes during liver regeneration (By similarity). Plays a role in the vascular response to injury (By similarity).

### **Cellular Location**

Nucleus. Cytoplasm, cytosol. Mitochondrion Note=Nuclear export to the cytosol is XPO1-mediated and positively regulated by IFI27 (PubMed:22427340). Translocation to the mitochondrion upon interaction with RXRA and upon the presence of 9-cis retinoic acid (PubMed:17761950).

#### **Tissue Location**

Fetal muscle and adult liver, brain and thyroid.

### NUR77 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### NUR77 Rabbit mAb - Images

