

PDLIM7 Rabbit mAb
Catalog # AP75886**Specification**

PDLIM7 Rabbit mAb - Product Information

Application	WB
Primary Accession	Q9NR12
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	49845

PDLIM7 Rabbit mAb - Additional Information**Gene ID** 9260**Other Names**
PDLIM7**Dilution**
WB~~1/500-1/1000**Format**
Liquid**PDLIM7 Rabbit mAb - Protein Information****Name** PDLIM7**Synonyms** ENIGMA**Function**

May function as a scaffold on which the coordinated assembly of proteins can occur. May play a role as an adapter that, via its PDZ domain, localizes LIM-binding proteins to actin filaments of both skeletal muscle and nonmuscle tissues. Involved in both of the two fundamental mechanisms of bone formation, direct bone formation (e.g. embryonic flat bones mandible and cranium), and endochondral bone formation (e.g. embryonic long bone development). Plays a role during fracture repair. Involved in BMP6 signaling pathway (By similarity).

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Note=Colocalizes with RET to the cell periphery and in some cytoskeletal components. Colocalizes with TPM2 near the Z line in muscle. Colocalizes with TBX4 and TBX5 to actin filaments (By similarity).

Tissue Location

Isoform 1 and isoform 2 are expressed ubiquitously, however, isoform 2 predominates in skeletal muscle, isoform 1 is more abundant in lung, spleen, leukocytes and fetal liver

PDLIM7 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PDLIM7 Rabbit mAb - Images

