

## Cullin 4A (CUL4A) Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7590a

#### Specification

# Cullin 4A (CUL4A) Antibody (N-term) - Product Information

Application	WB.E
Primary Accession	013619
Other Accession	O3TCH7
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	87680
Antigen Region	102-131

## Cullin 4A (CUL4A) Antibody (N-term) - Additional Information

Gene ID 8451

**Other Names** Cullin-4A, CUL-4A, CUL4A

**Target/Specificity** This Cullin 4A (CUL4A) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 102-131 amino acids from the N-terminal region of human Cullin 4A (CUL4A).

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** Cullin 4A (CUL4A) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Cullin 4A (CUL4A) Antibody (N-term) - Protein Information

Name CUL4A {ECO:0000303|PubMed:9721878, ECO:0000312|HGNC:HGNC:2554}



Function Core component of multiple cullin-RING-based E3 ubiquitin- protein ligase complexes which mediate the ubiquitination of target proteins (PubMed:14578910, PubMed:14739464, PubMed:15448697, PubMed:15548678, PubMed:15811626, PubMed:16678110, PubMed:17041588, PubMed:24209620, PubMed:30166453, PubMed:33854232, PubMed:<u>33854239</u>). As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme (PubMed: 14578910, PubMed: 14739464, PubMed:15448697, PubMed:15548678, PubMed:15811626, PubMed:16678110, PubMed: 17041588, PubMed: 24209620). The E3 ubiquitin- protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (PubMed: 14578910, PubMed: 14739464, PubMed:15448697, PubMed:15548678, PubMed:15811626, PubMed:16678110, PubMed:<u>17041588</u>, PubMed:<u>24209620</u>). The functional specificity of the E3 ubiguitin-protein ligase complex depends on the variable substrate recognition component (PubMed: 14578910, PubMed:14739464, PubMed:15448697, PubMed:15548678, PubMed:15811626, PubMed:16678110, PubMed:17041588, PubMed:24209620). DCX(DET1-COP1) directs ubiguitination of JUN (PubMed: 14739464). DCX(DDB2) directs ubiguitination of XPC (PubMed: 15811626). DCX(DDB2) ubiquitinates histones H3-H4 and is required for efficient histone deposition during replication-coupled (H3.1) and replication-independent (H3.3) nucleosome assembly, probably by facilitating the transfer of H3 from ASF1A/ASF1B to other chaperones involved in histone deposition (PubMed: 16678110, PubMed: 17041588, PubMed: 24209620). DCX(DTL) plays a role in PCNA-dependent polyubiguitination of CDT1 and MDM2-dependent ubiguitination of p53/TP53 in response to radiation-induced DNA damage and during DNA replication (PubMed:14578910, PubMed:15448697, PubMed:15548678). DCX(DTL) directs autoubiguitination of DTL (PubMed: 23478445). In association with DDB1 and SKP2 probably is involved in ubiguitination of CDKN1B/p27kip (PubMed: 16537899). Is involved in ubiguitination of HOXA9 (PubMed: 14609952). The DDB1-CUL4A- DTL E3 ligase complex regulates the circadian clock function by mediating the ubiquitination and degradation of CRY1 (PubMed: 26431207). The DCX(ERCC8) complex (also named CSA complex) plays a role in transcription-coupled repair (TCR) (PubMed:<u>12732143</u>, PubMed:<u>32355176</u>, PubMed:<u>38316879</u>). A number of DCX complexes (containing either TRPC4AP or DCAF12 as substrate-recognition component) are part of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiguitination and degradation (PubMed:29779948). The DCX(AMBRA1) complex is a master regulator of the transition from G1 to S cell phase by mediating ubiguitination of phosphorylated cyclin-D (CCND1, CCND2 and CCND3) (PubMed:<u>33854232</u>, PubMed:<u>33854239</u>). The DCX(AMBRA1) complex also acts as a regulator of Cul5-RING (CRL5) E3 ubiquitin-protein ligase complexes by mediating ubiquitination and degradation of Elongin-C (ELOC) component of CRL5 complexes (PubMed: 30166453). With CUL4B, contributes to ribosome biogenesis (PubMed: 26711351).

#### Cullin 4A (CUL4A) Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Cullin 4A (CUL4A) Antibody (N-term) - Images





Western blot analysis of CUL4a (arrow) using rabbit polyclonal CUL4a Antibody (Human N-term) (Cat.#AP7590a).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the CUL4a gene (Lane 2) (Origene Technologies).

# Cullin 4A (CUL4A) Antibody (N-term) - Background

CUL4a is a core component of multiple cullin-RING-based E3 ubiquitin-protein ligase complexes which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1. The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition component. DCX(DET1-COP1) directs ubiquitination of JUN. DCX(DDB2) directs ubiquitination of XPC. In association with RBX1, DDB1 and DDB2 is required for histone H3 and histone H4 ubiquitination in response to ultraviolet and may be important for subsequent DNA repair. DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitiantion of TP53 in response to radiation-induced DNA damage and during DNA replication. In association with DDB1 and SKP2 probably is involved in ubiquitination of CDKN1B/p27kip. Is involved in ubiquitination of HOXA9.

## Cullin 4A (CUL4A) Antibody (N-term) - References

Huang,J.,Oncogene 27 (29), 4056-4064 (2008) Hu,J.,Genes Dev. 22 (7), 866-871 (2008) Wen,X., J. Biol. Chem. 282 (37), 27046-27057 (2007)