

**SPI1 Rabbit mAb**  
**Catalog # AP75976****Specification**

---

**SPI1 Rabbit mAb - Product Information**

Application	<b>WB, IHC-P</b>
Primary Accession	<a href="#">P17947</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>31083</b>

**SPI1 Rabbit mAb - Additional Information****Gene ID** 6688**Other Names**

SPI1

**Dilution**

WB~~1/500-1/1000

IHC-P~~N/A

**Format**

Liquid

**SPI1 Rabbit mAb - Protein Information****Name** SPI1**Function**

Pioneer transcription factor, which controls hematopoietic cell fate by decompacting stem cell heterochromatin and allowing other transcription factors to enter otherwise inaccessible genomic sites. Once in open chromatin, can directly control gene expression by binding genetic regulatory elements and can also more broadly influence transcription by recruiting transcription factors, such as interferon regulatory factors (IRFs), to otherwise inaccessible genomic regions (PubMed:<a href="http://www.uniprot.org/citations/23658224" target="\_blank">23658224</a>, PubMed:<a href="http://www.uniprot.org/citations/33951726" target="\_blank">33951726</a>). Transcriptionally activates genes important for myeloid and lymphoid lineages, such as CSF1R (By similarity). Transcriptional activation from certain promoters, possibly containing low affinity binding sites, is achieved cooperatively with other transcription factors. FCER1A transactivation is achieved in cooperation with GATA1 (By similarity). May be particularly important for the pro- to pre-B cell transition (PubMed:<a href="http://www.uniprot.org/citations/33951726" target="\_blank">33951726</a>). Binds (via the ETS domain) onto the purine-rich DNA core sequence 5'-GAGGAA-3', also known as the PU-box (PubMed:<a href="http://www.uniprot.org/citations/33951726" target="\_blank">33951726</a>). In vitro can bind RNA and interfere with pre-mRNA splicing (By similarity).

**Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00237, ECO:0000269|PubMed:33951726}

**Tissue Location**

In the bone marrow, concentrated in hematopoietic stem cell, lymphoid progenitor, myeloid lineage (granulocyte macrophage progenitors, classical dendritic cells, monocytes) and B-cell clusters Among B-cells, predominantly expressed in pre-B1 cells (PubMed:33951726). Expressed in germinal center B-cells (PubMed:23166356).

**SPI1 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**SPI1 Rabbit mAb - Images**